

WEED MANAGEMENT IN LANDSCAPES

Integrated Pest Management for Landscape Professionals and Home Gardeners

Weed management in landscape plantings is often made difficult by the complexity of many plantings: usually more than one species is planted in the landscaped area and there is a mix of annual and perennial ornamentals. The great variety of ornamental species, soil types, slopes, and mulches creates the need for a variety of weed management options. There are also considerations regarding public concern about the use of chemicals to control weeds and their effect on water quality if the herbicide moves offsite through runoff. The choice of a specific weed management program depends on the weeds present and the types of turf or ornamentals planted in the area. Because of the many variables, weeds in landscape plantings are usually controlled by a combination of nonchemical and chemical methods.

Most landscape plantings include turfgrass, bedding plants, herbaceous perennials, shrubs, and trees. Use this publication as a practical review and guide to weed management options suited to planting beds and areas around trees and shrubs. Information on integrated pest management more directed towards turfgrass can be found in *Pest Notes: Weed Management in Lawns* (see References).

WEED MANAGEMENT BEFORE PLANTING

An integrated approach, utilizing several methods, is the most economical and effective means of controlling weeds. Develop your weed management plan for landscapes before you plant by following these five basic steps:

1. *Site assessment.* Before soil preparation and when the weeds are visible, evaluate the soil, mulch, and slope of

the site so problems can be corrected or future problems anticipated before planting. Site characteristics to look for include drainage, soil compaction, shading, and water infiltration rate. Identify the weed species in the area, with particular focus on perennial weeds (see References). The best time to look for winter annual weeds is mid-to late winter; perennials and summer annuals are easiest to identify in mid-to late summer.

2. *Site preparation.* The most often overlooked aspect of a landscape maintenance program is site preparation. Control existing weeds, especially perennials, before any grading and development are started. Glyphosate (Roundup, etc.) can be used to kill existing annual and perennial weeds. Preplant treatment with fumigants (available to licensed pesticide applicators only) or soil solarization can be used if time allows; however, 6 weeks are required for solarization and it is most effective when done during the time of highest sun radiation—from June to August in California (see References). Annual weeds can be reduced by irrigating the area after final grading, allowing the weeds to emerge. While the weeds are still small, a shallow cultivation (less than 1 inch), scraping the weeds off the soil, or spraying with a postmergent non-selective herbicide and then repeating this process of irrigation, emergence, and removal 2 or 3 times will greatly reduce annual weed competition and population.

3. *Define the type of planting.* There are more weed control options if the planting consists entirely of woody plants as opposed to herbaceous annuals or perennial plants, or a mixture of all three.

If there is opportunity to do so, consider using mowing strips, planter beds, and borders to reduce weed encroachment and delineate planting types (Fig. 1). Choosing drip irrigation and micro-sprinklers over conventional or flood sprinkler heads will also help in reducing weeds since the water is placed at or near your plants. Watering large areas where there are no plants will only encourage weeds to grow there.

4. *Don't introduce weeds.* Weeds are sometimes introduced in the soil brought to the landscape site either when amending the soil or in the potting mix of transplants.

5. *Encourage rapid establishment of desired plants.* Use the best management practices to get the plants established as quickly as possible so that they become competitive with weeds and more tolerant of herbicides applied to the site. Hand-weeding and keeping weeds from producing seeds in the landscape will greatly reduce overall weed populations.



Figure 1. Separate shrub areas from invasive lawns or ground covers by using headers. Headers are concrete, metal, plastic, or wood barriers extending 8 inches or more below ground and 2 to 3 inches above ground.

PEST NOTES

University of California
Agriculture and Natural Resources

Publication 7441

Revised March 2007

WEED MANAGEMENT AFTER PLANTING

When developing a weed management plan for an existing planting or after an installation is in place, consider the types of landscape plants present and the weeds present and their life cycles (annual, biennial, perennial) (Table 1).

Weed control options in the landscape include hand-weeding and cultivation, mowing, mulching, hot water treatments, and chemical control. All of these methods are used at one time or another in landscape maintenance operations (Table 2). For instance, after elimination by hand-pulling, cultivation, or a postmergent herbicide application, the subsequent growth of annual weeds can be discouraged with mulches and/or pre-emergent herbicides.

TABLE 1. Common Weeds in Landscape Plantings.

Annuals

- annual bluegrass
- clover (black medic and burclover)
- common groundsel +
- crabgrass (large and smooth) +
- little mallow (cheeseweed)
- pigweed (redroot and prostrate)
- prickly lettuce
- purslane
- sowthistle
- spurge (prostrate and creeping) +
- wild barley
- wild oat

Biennials

- bristly oxtongue

Perennials

- bermudagrass +
- creeping woodsorrel +
- dandelion
- field bindweed +
- kikuyugrass
- nutsedge (yellow and purple) +
- oxalis (creeping woodsorrel and Bermuda buttercup)

See the online weed photo gallery at www.ipm.ucdavis.edu for photos and information about these and many other weeds.

+ especially troublesome

Hand-weeding and Cultivation

Cultivation (hoeing) and hand-weeding selectively remove weeds from ornamental plantings. If weeds are scattered throughout the site, hand-weeding may be the preferred management method. Hand-weeding can be time consuming, but should be included in all weed management programs to keep weeds from seeding. Hand-weeding must be repeated frequently until the plantings become established. Cultivation can damage ornamentals with shallow roots, bring weed seeds to the soil surface, and propagate perennial weeds. When cultivating, avoid deep tilling, as this brings buried weed seeds to the soil surface where they are more likely to germinate.

Perennial weeds are often spread by cultivation and should be controlled or removed by other methods. Frequent removal of weeds when they are small and have not yet set seed will rapidly reduce the number of annual weeds.

Flaming. Young weeds in open areas also can be controlled with small flaming units. Propane burners are available to rapidly pass over young weeds to kill them. A quick pass over the plant is all that is necessary; do not burn the weed to the ground. Flaming is more effective on broadleaf weeds than grasses. Be careful not to flame dry vegetation, dry wood chips, or near buildings and other flammable materials, and don't get the flame near desired plants.

Mowing

Mowing can be used to prevent the formation and spread of weed seeds from many broadleaf weeds into cultivated areas by cutting off flower heads. However, weeds that flower lower than the mowing blade (such as spotted spurge or common woodsorrel) are not controlled. Repeated mowing tends to favor the establishment of grasses and low-growing perennial weeds. Mowing of some ground covers can rejuvenate them and make them more competitive against weeds.

String Trimmers. The top growth of older weeds can be controlled by using

a string trimmer. Annual broadleaf weeds are more effectively controlled than annual grasses because the growing points of grasses are usually below ground. Most perennial weeds are not controlled using string trimmers but will make them less noticeable in areas if you are limited in other methods for control.

Prevent repeated damage to trunks of established woody perennials from the string trimmer by placing a temporary shield around the trunk. Create your own shield by placing a segment of irrigation pipe that is several inches in diameter and split lengthwise around the base of the trunk.

Mulches

A mulch is any material placed on the soil to cover and protect it. Mulches suppress annual weeds by limiting light required for weed establishment. Many types of landscape mulches are available. The most common are bark and other wood products and black plastic or landscape fabric materials. Other products that are used include paper, yard compost, hulls from nuts (such as almonds) or cereals (rice), municipal composts, and stones.

Organic Mulches. Organic mulches include wood chips, sawdust, yard waste (leaves, clippings, and wood products), and hardwood or softwood bark chips or nuggets. Bark chips are moderate-sized particles (¼ to ½ inch) and have moderate to good ability to withstand decomposition, while bark nuggets are larger in size (½ to 1½ inches) and have excellent stability over time. All of these can be used in landscape beds planted with herbaceous or woody ornamentals. Larger mulch pieces (greater than 1½ inch) do not provide good weed control the space between the pieces allows weeds to grow through.

The thickness or depth of a mulch necessary to adequately suppress weed growth depends on the mulch type and the weed pressure. The larger the particle size of the organic mulch, the greater the depth required to exclude all light from the soil surface. Coarse-

textured mulches can be applied up to 4 inches deep and provide long-term weed control. Fine-textured mulches pack more tightly and should only be applied to a depth of about 2 inches. If the mulch is too decomposed, it is a weed propagation medium rather than a means of prevention. Plan to periodically replenish organic mulches, regardless of particle size, because of

decomposition, movement, or settling. If seedlings germinate in mulches, a light raking, hoeing, or hand-weeding will remove the young weeds.

Inorganic Mulches. Including both natural and synthetic products, these mulches are generally more expensive and less widely used in the landscape. Natural inorganic mulches are stable

over time and include materials such as sand, gravel, or pebbles. Most of these products are used in public and commercial plantings. If using a rock mulch, consider placing a landscape fabric underneath it. The fabric creates a layer between the mulch and soil, preventing rock pieces from sinking into the soil. The fabric prevents soil from moving above the rock layer,

TABLE 2. How to Manage Weeds in Five Types of Landscape Plantings.

Type of planting and comments	Recommendations
<p>Woody Trees and Shrub Beds. Densely shaded plantings reduce weeds. Preplant weed control is not as critical as in other types of plantings. It is often necessary to combine treatments for complete weed control.</p>	<p>Control perennial weeds before planting (although control may be possible after planting); use geotextile fabrics with a shallow layer of mulch or use a thick layer of mulch without a geotextile base; use a preemergent herbicide, if needed, and supplement with spot applications of postemergent herbicides and/or hand-weeding. Perennial weeds may be controlled by manual removal, spot applications of glyphosate or glufosinate, or, in some instances, dormant-season applications of preemergent herbicides. Escaped weeds may be controlled manually or with spot applications of postemergent herbicides.</p>
<p>Woody Ground Cover Beds. Woody mature ground covers should exclude most weeds; however, weed encroachment during establishment is likely.</p>	<p>Control perennial weeds before planting, although perennial grasses may be selectively controlled after planting with fluzifop (Fusilade,¹ Ornamec¹), clethodim (Envoy¹), sethoxydim (Grass-Getter), or other selective grass herbicides. Annual weeds may be controlled with mulch plus a preemergent herbicide, supplemented with some hand-weeding. Use geotextiles where possible but do not use them where ground covers are expected to root and spread. After planting, it is difficult to make spot applications of nonselective herbicides without injuring desirable plants. Postemergent control of most annual and perennial grasses is possible.</p>
<p>Annual Flower Beds. A closed canopy will help shade out many weeds. Periodic cultivations (at 3- to 4-week intervals and between display rotations) will suppress many weeds.</p>	<p>Control perennial weeds before planting and carefully select flower species for weed management compatibility. Annual weeds may be controlled with mulches, preemergent herbicides, frequent cultivation, and/or hand-weeding. Perennial grasses can be selectively controlled with clethodim or fluzifop, or other grass-selective herbicides, but other perennial weeds cannot be selectively controlled after planting. Geotextiles generally are not useful because of the short-term nature of the planting. Avoid nonselective herbicides after planting.</p>
<p>Herbaceous Perennial Beds. Weed management options in herbaceous perennial beds are similar to those for annual flowers, except (1) it is more important to eradicate perennial weeds as there will be no opportunity to cultivate or renovate the bed for several years; and (2) fewer species are included on herbicide labels.</p>	<p>Control perennial weeds before planting; use geotextiles where possible; use mulches with a preemergent herbicide; and supplement with hand-weeding.</p>
<p>Mixed Plantings of Woody and Herbaceous Plants. Weed management is complex because of the diversity of species. Different areas of the bed could receive different treatments. Site preparation is critical because postplant herbicide choices are few.</p>	<p>Plant the woody species first; control perennial weeds in the first two growing seasons, then introduce the herbaceous species. Plant close together to shade the entire area. Another option may be to define use-areas within the bed that will receive similar weed management programs.</p>

Herbicides should not normally be needed by home gardeners. Mulching, removal by hand, and proper placement of irrigation water will be sufficient in most cases.

¹For use only by professional applicators.

which would bring weed seed to the surface.

Black plastic (solid polyethylene) can be used underneath mulches to improve weed control. It provides excellent control of annual weeds and suppresses perennial weeds, but lacks porosity and restricts air and water movement. It also tends to tear and break apart rather quickly. For this reason, black plastic may not be the preferred long-term weed control method in landscape beds.

Synthetic Mulches. Manufactured materials, called geotextiles or landscape fabrics, have been developed to replace black plastic in the landscape. Geotextiles are porous and allow water and air to pass through them, overcoming the major disadvantage of black plastic. Although these materials are relatively expensive and time-consuming to install, they become cost-effective if the planting is to remain in place for 4 or more years. Geotextiles are used mainly for long-term weed control in woody ornamental trees and shrubs. Geotextiles should not be used where the area is to be replanted periodically, such as in annual flower beds or in areas where the fabric would inhibit the rooting and spread of ground covers. Tree and shrub roots can penetrate the materials and if the material is removed, damage can occur to the plant's root system. This might be a concern if a fabric has been in place longer than 5 years. One geotextile fabric (BioBarrier) has an herbicide encapsulated in nodules on the fabric that reduces root penetration problems.

Placing a landscape fabric under mulch results in greater weed control than mulch used alone. There are differences in the weed-controlling ability among the geotextiles: fabrics that are thin, lightweight, or have an open mesh allow for greater weed penetration than more closely woven or nonwoven (spunbonded) fabrics.

To install a landscape fabric, you can plant first and then install the fabric afterwards using U-shaped nails to peg

it down. After laying the cloth close to the ground, cut an "X" over the plant and pull it through the cloth. If laying down a fabric before planting, cut an "X" through the fabric and dig a planting hole. Avoid leaving soil from the planting hole on top of the fabric because this could put weed seeds above the material. Fold the "X" back down to keep the geotextile sheet as continuous as possible. Weeds will grow through any gap in the landscape fabric, so it is important to overlap pieces of fabric and tack them down tightly. Apply a shallow mulch layer (about 1 inch deep) to thoroughly cover the fabric and prevent photodegradation. If weeds grow into or through the geotextile, remove them when they are small to prevent them from creating holes in the fabric. Maintain a weed-free mulch layer on top of the fabric by hand-weeding or by applying herbicides. Use of a rock mulch above a landscape fabric can have greater weed control than fabric plus organic mulch combinations, but may stress plants in some situations. See Problems section following.

Special consideration should be given to the fact that yellow nutsedge grows through all geotextiles. However, some fabrics are better at suppressing yellow nutsedge than others. For more information, see *Pest Notes: Nutsedge* listed in References.

Problems with Organic and Natural Inorganic Mulches. There are several problems associated with the use of organic and inorganic mulches. Perennial weeds such as field bindweed and nutsedges often have sufficient root reserves to enable them to penetrate even thick layers of mulches. Some annual weeds will grow through mulches, while others may germinate on top of the mulch as it decomposes. Weeds that are a particular problem are those that have windborne seeds such as common groundsel, prickly lettuce, and common sowthistle. Applying mulches at depths of greater than 4 inches may injure plants by keeping the soil too wet and limiting oxygen to the plant's roots. Disease

incidence, such as root or stem rot, may increase when deep mulches are maintained.

When mulches are too fine, applied too thickly, or begin to decompose, they stay wet between rains and allow weeds to germinate and grow directly in the mulch. For best weed control, use a coarse-textured mulch with a low water-holding capacity. When used alone, mulches rarely provide 100% weed control. To improve the level of weed control, one can apply preemergent herbicides at the same time as the mulch (see Table 3). Supplemental hand-weeding or spot spraying may also be needed.

Avoid mulches with a pH less than 4 or that have an "off odor" such as ammonia, vinegar, or rotten egg smell. These mulches were stored incorrectly and contain chemical compounds that may injure plants, especially herbaceous plants.

If using a composted mulch, temperatures achieved during the composting process should have killed most weed seeds. However, if the compost was stored uncovered in the open, weed seeds may have been blown onto the mulch. Be sure the mulch is not contaminated with weed seeds or other propagules such as nutsedge tubers.

Rock mulches should always be used with a landscape fabric underneath. Avoid getting soil or other growing media between the rocks or else weeds will grow there. Removal of these weeds by hand weeding or hoeing is very difficult. White rocks are very reflective and this increased light can damage sensitive plants. Dark colored rocks will retain heat and may also cause plant stress.

Hot Water or Steam Treatments

There are several machines currently available that use superheated hot water or steam to kill weeds. The equipment is expensive to purchase and maintain, so these machines are not appropriate for home use. However, commercial landscapers may find them useful in certain situations where the

use of herbicides is not desired such as when line-marking playing fields, in playgrounds, around woody plants, for edging, and for weeds growing along fence lines. Some brands of equipment travel slowly (about 2 mile/hour) and are probably not cost-effective for weed control along roadsides. Because these methods employ boiling water or steam, workers must be adequately trained in the use of the machines to prevent severe burns. These machines are most effective on very young annual weeds or perennials that have recently emerged from seeds. The effect is similar to that of a nonselective, postemergent herbicide. Hot water and steam are not very effective on perennial weeds with established storage organs, such as rhizomes and bulbs, nor do they control woody plants. In general, broadleaf weeds are more easily controlled by this method than grasses. A home gardener can apply hot water (greater than 200° F) and achieve a limited amount of weed damage. This method is not very effective because the water temperature rapidly drops once it is removed from the heat source.

Herbicides for Landscape Plantings

Herbicides have been effectively used in many types of landscape plantings and are most often integrated with the cultural practices discussed above. Generally, home gardeners should not need to apply herbicides to existing landscape plantings. Hand-weeding and mulching should provide sufficient control and avoid hazards to desirable plants associated with herbicide use. Many herbicides listed here are for use by professional landscape pest managers and are not available to home gardeners. To determine which herbicide(s) are in a product, read the active ingredients information on the label.

Preemergent Herbicides. When weeds have been removed from an area, preemergent herbicides can then be applied to prevent the germination or survival of weed seedlings. Preemergent herbicides must be applied before the weed seedlings emerge. Examples

of preemergent herbicides include: DCPA (Dacthal), dithiopyr (Dimension), isoxaben (Gallery), napropamide (Devrinol), oryzalin (Surflan, Weed Impede), oxadiazon (Ronstar), oxyfluorfen (Goal), pendimethalin (Pendulum, Pre-M), and prodiamine (Barricade). DCPA, dithiopyr, oryzalin, napropamide, pendimethalin, and prodiamine control annual grasses and many broadleaf weeds and can be used safely around many woody and herbaceous ornamentals. Isoxaben is used for control of broadleaf weeds.

Match herbicides with weeds present, and consider using herbicide combinations. Combinations of herbicides increase the spectrum of weeds controlled and provide effective control of grasses and many broadleaf weeds. Commonly used combinations include tank mixes of the materials listed above or isoxaben/trifluralin (Snapshot), oryzalin/benfen (XL2G), oxyfluorfen/oryzalin (Rout), and oxyfluorfen/pendimethalin (Ornamental Herbicide 2). Check the label to determine which ornamental species the material can safely be used around and which species of weeds are controlled.

Timing. Timing of a preemergent herbicide application is determined by when the target weed germinates, or by when the weed is in the stage that is most sensitive to the herbicide. In general, late summer/early fall applications of preemergent herbicides are used to control winter annuals, while late winter/early spring applications are used to control summer annuals and seedlings of perennial weeds. If heavy rainfall occurs after preemergent herbicide application or if a short residual (short acting) product was applied, a second preemergent herbicide application may be needed. Generally, herbicides degrade faster under wet, warm conditions than under dry, cool conditions. Some preemergent herbicides, such as surflan, need to be applied at a higher rate if the soil is high in organic matter due to binding with the organic matter. Check the herbicide label to recommendations about what rate to use depending on your soil type.

Cultivation and Preemergent Herbicides. No cultivation should occur after an application of oxyfluorfen; however, shallow cultivation (1 to 2 inches) will not affect the activity of napropamide, pendimethalin, or oryzalin. Also, soil type and pH can affect the activity of some herbicides. Use the information contained in herbicide labels and from your local county Cooperative Extension office to determine the tolerance of an ornamental plant species to a given herbicide.

Herbicide and Mulch Placement. The placement of a preemergent herbicide in relation to an organic mulch can affect the herbicide's performance. Additionally, the characteristics of organic mulches can affect how herbicides work. Mulch that is made up of coarse particles will have little effect on herbicide activity. A mulch that primarily consists of fine particles can reduce the availability of some herbicides. The finer the organic material (compost or manure, compared to bark), the greater the binding of the herbicide. Many preemergent herbicides are tightly bound by organic matter, and while the binding minimizes leaching, it can also minimize an herbicide's activity.

Another important factor is the depth of the mulch. An herbicide applied on top of a thin mulch may be able to leach through to where the weed seeds are germinating, but when applied to the top of a thick layer of mulch it may not get down to the zone of weed seed germination. Products like oxadiazon (Ronstar) and oxyfluorfen (Goal) that require a continuous surface layer must be placed on the soil surface under the mulch. Suggestions for use of mulch with herbicides are given in Table 3.

Postmergent Herbicides. When weeds escape preemergent herbicides or geotextile fabrics, postmergent herbicides can be used to control established weeds. Postmergent herbicides control existing plants only and do not give residual weed control. Their primary function is to control young annual species, but they are also used to control perennial species. Some herbicides are systemic, that is, they

are translocated throughout rapidly growing plants and can kill the entire plant. Common systemic herbicides include herbicides clethodim and fluazifop which selectively control most annual and perennial grasses, 2, 4-D which selectively controls many annual and perennial broadleaf plants and glyphosate (Roundup Pro and others) which is nonselective and effective on most annual and perennial weeds. Glufosinate (Finale), diquat (Reward), pelargonic acid (Scythe), and the new herbicides containing clove oil (Matran, EcoEXEMPT) are nonselective, contact herbicides that kill or injure any vegetation they contact. They kill small annual weeds, but only "burn off" the tops of perennial weeds.

Most of the specific herbicides listed here are not available for home gardener use and should only be applied by professionals. The exceptions are dithiopyr (in Greenlight Crabgrass Preventer, some formulations of Preen, and some "Weed and Feed" mixes); fluazifop (in Ortho Grass B Gon, Greenlight Grass Out, and others) glufosinate (Finale and others), pelargonic acid (Scythe), clove oil (Burnout and others), and glyphosate (Roundup and others).

Avoid Herbicide Injury. Because of the close proximity of many different species of plants in the landscape, herbicide injury may occur, resulting in visual plant damage. Herbicide injury symptoms vary according to plant species and the herbicide and can include the following.

- yellowing (chlorosis)
- bleaching
- root stunting
- distorted growth
- death of leaves

Granular formulations of preemergent herbicides are less likely to cause injury than sprayable formulations. Using a granular formulation reduces the potential for damage by foliar uptake, but granules of oxadiazon (Ronstar) or oxyfluorfen (Goal) mixtures can still injure plants if they collect in the base of leaves or adhere to wet leaves. Apply nonselective liquid herbicides such as diquat, pelargonic acid, or glyphosate

TABLE 3. Suggestions for Placement of Herbicide with an Organic Mulch.

Herbicide	Application
Devrinol (napropamide)	under the mulch
Dimension (dithiopyr)	best under mulch, but provides some control when applied on top. Limited use on some landscape species.
Gallery (isoxaben)	best under the mulch, moderate control when applied on top of mulch
Ornamental Herbicide 2 (pendimethalin plus oxyfluorfen)	works well both under or over mulch
Pennant (metolachlor)	under the mulch
Ronstar (oxadiazon)	over the mulch
Rout (oryzalin plus oxyfluorfen)	works well both under or over mulch
Surflan (oryzalin)	best under the mulch but provides some control when applied on top of mulch
Surflan plus Gallery	under the mulch but will give a fair amount of control even when applied on top of mulch
Treflan (trifluralin)	under the mulch
XL2G (oryzalin/benfen)	under the mulch

with low pressure and large droplets on a calm day. Use shielded sprayers when making applications around ornamentals to avoid contact with non-target plants.

Herbicide injury to established plants from soil-applied chemicals is often temporary but can cause serious growth inhibition to newly planted ornamentals. Herbicides that contain oryzalin or isoxaben are more likely to cause this injury. Injury may result when persistent herbicides are applied to surrounding areas for weed control in turf, agronomic crops, or complete vegetative control under pavement. Activated charcoal incorporated into the soil may adsorb the herbicide and minimize injury. Usually it just takes time for herbicide residues to completely degrade. To speed degradation, supplement the organic content of the soil and keep it moist but not wet during periods of warm weather.

REFERENCES

Derr, J. F. et al. Feb. 1997. Weed Management in Landscape and Nursery Plantings, from Weed Management and Horticultural Crops. WSSA/ASHS Symposium.

DiTomaso, J. M. and E. A. Healy. 2007. *Weeds of California and Other Western States*. Oakland: Univ. Calif. Agric Nat. Res. Publ. 3488

Dreistadt, S. H. 1992. *Pests of Landscape Trees and Shrubs*. Oakland: Univ. Calif. Agric Nat. Res. Publ. 3359.

Elmore, C.L. 1997. *Soil Solarization: A Nonpesticidal Method for Controlling Diseases, Nematodes, and Weeds*. Oakland: Univ. Calif. Agric. Nat. Res. Publication 21377

Fischer, B. B., ed. 1998. *Grower's Weed Identification Handbook*. Oakland: Univ. Calif. Agric Nat. Res. Publ. 4030.

UC Statewide IPM Program. Integrated Pest Management Weed Photo Gallery, an online photographic reference to many weeds commonly found in California. Available online, http://ipm.ucdavis.edu/PMG/weeds_common.html. Accessed Jan. 16, 2007.

UC Statewide IPM Program. The *Pest Notes* series has many relevant titles: *Annual Bluegrass*. *Bermudagrass*. *Common Knotweed*. *Common Purslane*. *Crabgrass*. *Creeping Woodsorrel/Bermuda Buttercup*. *Dandelion*. *Dodder*. *Field*

Bindweed. Green Kyllinga. Kikuyugrass. Mistletoe. Nutsedge. Poison Oak. Plantains. Russian Thistle. Spotted Spurge. Weed Management in Lawns. Wild Blackberries. Oakland: Univ. Calif. Agric Nat. Res. Available online, <http://www.ipm.ucdavis.edu/PMG/menu.homegarden.html>.

UC Statewide IPM Program. Soil solarization method described online, <http://ipm.ucdavis.edu/TOOLS/TURF/SITEPREP/soilsolar.html>. Accessed Jan. 16, 2007

Whitson, T.D. ed. 2004. *Weeds of the West*. Western Society of Weed Science. ❖

For more information contact the University of California Cooperative Extension in your county. See your telephone directory for addresses and phone numbers.

AUTHORS: C. A. Wilen, UC Statewide IPM Program, San Diego Co.; C. L. Elmore, Weed Science/Vegetable Crops Emeritus, UC Davis

TECHNICAL EDITOR: M. L. Flint

DESIGN AND PRODUCTION: P. N. Galin

ILLUSTRATIONS: **Fig. 1** and **Fig. 2** Adapted from Dreistadt, S. H. 1992. *Pests of Landscape Trees and Shrubs*. Oakland: Univ. Calif. Agric Nat. Res. Publ. 3359.

Produced by IPM Education and Publications, UC Statewide IPM Project, University of California, Davis, CA 95616-8620

This Pest Note is available on the World Wide Web (www.ipm.ucdavis.edu)



This publication has been anonymously peer reviewed for technical accuracy by University of California scientists and other qualified professionals. This review process was managed by the ANR Associate Editor for Pest Management.

To simplify information, trade names of products have been used. No endorsement of named products is intended, nor is criticism implied of similar products that are not mentioned.

This material is partially based upon work supported by the Extension Service, U.S. Department of Agriculture, under special project Section 3(d), Integrated Pest Management.

WARNING ON THE USE OF CHEMICALS

Pesticides are poisonous. Always read and carefully follow all precautions and safety recommendations given on the container label. Store all chemicals in the original labeled containers in a locked cabinet or shed, away from food or feeds, and out of the reach of children, unauthorized persons, pets, and livestock.

Pesticides applied in your home and landscape can move and contaminate creeks, lakes, and rivers. Confine chemicals to the property being treated. Avoid drift onto neighboring properties, especially gardens containing fruits or vegetables ready to be picked.

Do not place containers containing pesticide in the trash or pour pesticides down sink or toilet. Either use the pesticide according to the label or take unwanted pesticides to a Household Hazardous Waste Collection site. Contact your county agricultural commissioner for additional information on safe container disposal and for the location of the Household Hazardous Waste Collection site nearest you. Dispose of empty containers by following label directions. Never reuse or burn the containers or dispose of them in such a manner that they may contaminate water supplies or natural waterways.

The University of California prohibits discrimination or harassment of any person on the basis of race, color, national origin, religion, sex, gender identity, pregnancy (including childbirth, and medical conditions related to pregnancy or childbirth), physical or mental disability, medical condition (cancer-related or genetic characteristics), ancestry, marital status, age, sexual orientation, citizenship, or status as a covered veteran (covered veterans are special disabled veterans, recently separated veterans, Vietnam era veterans, or any other veterans who served on active duty during a war or in a campaign or expedition for which a campaign badge has been authorized) in any of its programs or activities. University policy is intended to be consistent with the provisions of applicable State and Federal laws. Inquiries regarding the University's nondiscrimination policies may be directed to the Affirmative Action/Staff Personnel Services Director, University of California, Agriculture and Natural Resources, 300 Lakeside Drive, 6th Floor, Oakland, CA 94612-3550, (510) 987-0096.