

Maine Leading Causes of Death: Ten Most Common Causes 2018

Maine CDC
Office of Data, Research and Vital Statistics



Revised version: March 1, 2021

Acknowledgments

This report serves as an update to the 2015 Maine Mortality report which was co-authored by Sara Huston, Finn Teach, and Mary Caron in 2018. Many of the formats, tables, and charts build off that work and update information in that report.

Carolyn Bancroft prepared this report. We would like to thank the following people for their contributions of data preparation, editing, graphic support and review: Anne Rogers, Kim Haggan, Finn Teach, and Sara Huston.

Executive Summary

At the beginning of the 21st century, heart disease was the leading cause of death among Maine residents. In 2002, cancer took over as the leading cause of death and has remained there for 16 years.

Together, heart disease and cancer were responsible for 49% of all Maine resident deaths in 2018. When combined with the third most common cause of death, unintentional injuries, the top three causes accounted for 55% of Maine deaths in 2018.

The age-adjusted all-cause mortality rate decreased from 1999-2018, though the mortality rate has plateaued since 2011 with no significant decrease or increase in the mortality rate since then. Unintentional injuries moved from the sixth leading cause of death in 2004 to the third leading cause of death in 2016. The rise in unintentional injury deaths was due primarily to drug related deaths and falls.

This report, along with knowledge of best practices in prevention and public health, may support Maine public health organizations to develop initiatives that help Maine people live longer and healthier lives.

Table of Contents

Acknowledgments and Executive Summary.....	2
Section 1: Leading Causes of Death in Maine,2018.....	4
Section 2: At-a-Glance Sheets.....	10
Maine vs. U.S.....	11
Males vs. Females	12
Trends over Time	13
Counties and Public Health Districts.....	14
Age Groups	16
Section 3: Cause-specific Figures	17
Section 4: Cause-specific Maps	23
Section 5: Tables	26
Technical Notes	96
References	98

Section 1:
Leading Causes of Death in Maine,
2018

Summary of key findings

Leading cause of death rankings

Key findings in comparison with the US

- **Cancer** is the leading cause of death in Maine and **Heart Disease** is the leading cause of death in the US (see Section 5, Table 1).

Key findings by Sex

- **Cancer** is the leading cause of death among both males and females in Maine followed by **Heart Disease** as the 2nd leading cause of death.
- **Unintentional injury** is the 3rd leading cause of death among males and the 7th leading cause of death among females.
- **Suicide** is the 7th leading cause of death among males and the 14th leading cause of death among females (see Section 5, Table 2).

Key findings by Age group

- **Unintentional injury** is the leading cause of death among those ages 1 to 44 followed by **suicide**.
- Among those ages 45 and older, **Cancer** is the leading cause of death followed by **Heart Disease** (see Section 5, Table 3).

Key findings by Race and Ethnicity

- The top two leading causes of death are **Cancer** and **Heart Disease** for all race/ethnicity groups in Maine (see Section 5, Tables 4-5).

Key findings by County of Residence:

- **Cancer** is the leading cause of death in all Maine counties except for Aroostook, Lincoln and Washington where **Heart Disease** is the leading cause of death (See Section 5, Table 6).

Key findings trends over time


- **Heart Disease** and **Cancer** changed places as the number one cause of death in Maine in 2002 with Cancer being the number 1 and Heart Disease the number 2 cause of death for the past 16 years (2002-2018).
- **Unintentional Injury** deaths vacillated between the 6th and 7th leading cause of death between 2000 and 2005, moving to the 4th leading cause by 2013 to the 3rd leading cause 2016-2018.
- **Cerebrovascular Disease** moved from the 3rd leading cause in 2000 to the 5th leading cause in 2013, where it remains as the 5th leading cause of death in 2018 (see Section 5, Table 2).

Mortality rates:

- The statewide age-adjusted mortality rate decreased from 865.9 per 100,000 in 1999 to 753.9 per 100,000 in 2018.
 - However, while there was an average annual decrease of -1.49% in mortality rate from 1999 to 2011, the decline in mortality plateaued starting in 2011 with no significant increase nor decrease since then (see Section 5, Table 8).
- Cancer and unintentional injury are the leading cause of years of potential life lost before age 75. Heart disease and suicide are additional leading causes of years of potential life lost (see Section 5, Table 7).
- The age-adjusted cancer mortality rate in Maine decreased over the past two decades from 214.6 per 100,000 in 1999 to 162.0 per 100,000 in 2018.
 - There is a significant downward trend in cancer mortality in Maine with an average annual percent change of -1.44%.
 - Cancer remains the leading cause of death in Maine. In 2018, Maine's cancer rate is higher than the US which was 149.1 per 100,000 in 2018 (see Section 5, Table 13).
- The age-adjusted heart disease mortality rate decreased from 238.7 per 100,000 in 1999 to 146.9 per 100,000 in 2018.
 - While there was an average annual decrease of -4.17% in heart disease mortality rate from 1999 to 2010, the decline in mortality plateaued starting in 2010 with no significant increase nor decrease (see Section, 5, Table 18).
- The age-adjusted unintentional injury mortality rate nearly doubled over the past two decades from 34.9 per 100,000 in 1999 to 64.7 per 100,000 in 2018.
 - Unintentional injury deaths in Maine have increased significantly since 2011, an average annual percent change of 11.13%.
 - Though the mortality rate due to unintentional injuries is higher among males compared with females, the rate has increased significantly for both sexes over the past 5 years (see Section 5, Tables 28-29).

For information about analysis of trends in mortality rates over time, please refer to the description of Joinpoint analysis in the Technical Notes section at the end of this report.

Top 10 Leading Causes of Death in Maine, 2018




Data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. October 2019.

In 2018, there were 14,716 deaths to Maine residents. Maine's leading cause of death was cancer. Other major causes included heart disease, unintentional injuries, chronic lower respiratory disease and cerebrovascular diseases.

Maine Leading Cause of Death Rank, 1999-2018

Unintentional injuries have risen from the 5th to the 3rd leading cause of death in Maine since 1999.




1999-2017 Data accessed on WONDER, US CDC. <https://wonder.cdc.gov/ucd-icd10.html>.

2018 data: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. October 2019.

Technical note: Ranks are based on number of deaths assigned to a rankable cause, not crude or age-adjusted rate comparisons, based on National Center for Health Statistics guidance.

Top 10 Leading Causes of Death, Maine vs. U.S., 2018




Maine data source: Maine CDC, Data Research, and Vital Statistics. October 2019.


US data source: US Centers for Disease Control and Prevention, National Center for Health Statistics. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Feb 13, 2020

Maine's death rates due to cancer, unintentional injury, chronic lower respiratory disease and suicide were significantly higher compared with the US rates in 2018.

Top 5 Leading Causes of Death in Maine, Ages 1-44, 2018




Top 5 Leading Causes of Death in Maine, Ages 45+, 2018



Data source: Maine CDC, Data Research, and Vital Statistics. January 2020.

Cancer was the leading cause of death for individuals over age 45 and unintentional injury was the leading cause of death among ages 1-44.


Leading Causes of Years of Potential Life Lost (YPLL) before Age 75, Maine, 2018



Data source: National Center for Health Statistics (NCHS), National Vital Statistics System. Centers for Disease Control and Prevention: WISQARS Leading Causes of Death. Available at: www.cdc.gov/injury/wisqars

Cancer and unintentional injury were the leading cause of years of potential life lost before age 75. Heart disease and suicide were other leading causes of years of potential life lost.

Preventable Causes of Death, Maine, 2018



Data source: Mokdad, Ali H., et al. "Actual causes of death in the United States, 2000." *JAMA* 291.10 (2004): 1238-1245.

Tobacco, poor diet and physical inactivity were the leading preventable causes of death in Maine. *Technical note:* These estimates are based on national research about what percentage of deaths are due to these preventable causes and they do not account for the opioid epidemic and thus illicit drug use is likely underestimated. The total number of Maine deaths for 2018 were multiplied by each percentage to produce the estimates above.

Section 2:

At-a-Glance Sheets

Maine vs. U.S. At-a-Glance Leading Causes of Death, 2018



Indicator	Unit	Maine	U.S.	Comparison	Year
All Deaths	Rate per 100,000	753.9	723.6	◆	2018
	95% CI	741.6 - 766.1	722.8 - 724.5		
Cancer Deaths	Rate per 100,000	162	149.1	◆	2018
	95% CI	156.3 - 167.6	148.7 - 149.5		
Heart Disease Deaths	Rate per 100,000	146.9	163.6	■	2018
	95% CI	141.6 - 152.3	163.2 - 164.0		
Unintentional Injury Deaths	Rate per 100,000	64.7	48	◆	2018
	95% CI	60.4 - 69.0	47.8 - 48.2		
Chronic Lower Respiratory Disease Deaths	Rate per 100,000	45.6	39.7	◆	2018
	95% CI	42.6 - 48.5	39.5 - 39.9		
Cerebrovascular Disease Deaths	Rate per 100,000	33.8	37.1	■	2018
	95% CI	31.2 - 36.3	36.9 - 37.3		
Alzheimer's Disease Deaths	Rate per 100,000	28.4	30.5	●	2018
	95% CI	26.0 - 30.7	30.3 - 30.7		
Diabetes Deaths	Rate per 100,000	23.4	21.4	●	2018
	95% CI	21.3 - 25.6	21.2 - 21.5		
Influenza and Pneumonia Deaths	Rate per 100,000	15.2	14.9	●	2018
	95% CI	13.5 - 16.9	14.8 - 15.0		
Suicide Deaths	Rate per 100,000	18.4	14.2	◆	2018
	95% CI	16.1 - 20.7	14.1 - 14.3		
Nephritis, Nephrotic Syndrome, and Nephrosis Deaths	Rate per 100,000	12.5	12.9	●	2018
	95% CI	11.0 - 14.1	12.8 - 13.0		

- ME doing better than the U.S.
 - ME similar to the U.S.
 - ◆ ME doing worse than the U.S.
- Rates are age-adjusted to the U.S. 2000 Standard Population.
95% CI: 95% Confidence Interval of the rate.

Maine data: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

US Data: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Feb 13, 2020.

February 2020. Carolyn Bancroft, Maine Center for Disease Control and Prevention.

The Department of Health and Human Services complies with applicable Federal and State civil rights laws and does not discriminate on the basis of disability, race, color, sex, gender, sexual orientation, age, national origin, religious or political belief, ancestry, familial or marital status, genetic information, association, previous assertion of a claim or right, or whistleblower activity, in admission or access to, or the operation of its policies, programs, services, or activities, or in hiring or employment practices.

Males vs. Females At-a-Glance

Leading Causes of Death, Maine, 2018



Indicator	Unit	Males	Females	Maine Total	Bar Chart	Year
All Deaths	Rate per 100,000	896.7	630	753.9		2018
	95% CI	876.5 - 917.0	615.1 - 644.8	741.6 - 766.1		
Cancer Deaths	Rate per 100,000	188.8	141.1	162.0		2018
	95% CI	179.7 - 198.0	133.9 - 148.4	156.3 - 167.6		
Heart Disease Deaths	Rate per 100,000	193.7	110.0	146.9		2018
	95% CI	184.2 - 203.2	103.9 - 116.1	141.6 - 152.3		
Unintentional Injury Deaths	Rate per 100,000	89.8	40.7	64.7		2018
	95% CI	82.4 - 97.3	36.3 - 45.2	60.4 - 69.0		
Chronic Lower Respiratory Disease Deaths	Rate per 100,000	53.5	39.7	45.5		2018
	95% CI	48.6 - 58.4	36.0 - 43.3	42.6 - 48.5		
Cerebrovascular Disease Deaths	Rate per 100,000	32.0	34.1	33.8		2018
	95% CI	28.1 - 35.8	30.7 - 37.4	31.2 - 36.3		
Alzheimer's Disease Deaths	Rate per 100,000	20.4	33.2	28.4		2018
	95% CI	17.3 - 23.5	29.9 - 36.4	26.0 - 30.7		
Diabetes Deaths	Rate per 100,000	29.4	18.1	23.4		2018
	95% CI	25.8 - 33.0	15.6 - 20.7	21.3 - 25.6		
Influenza and Pneumonia Deaths	Rate per 100,000	20.1	11.7	15.2		2018
	95% CI	16.9 - 23.2	9.8 - 13.7	13.5 - 16.9		
Suicide Deaths	Rate per 100,000	30.4	7.1	18.4		2018
	95% CI	26.1 - 34.6	5.2 - 9.5	16.1 - 20.7		
Nephritis, Nephrotic Syndrome and Nephrosis Deaths	Rate per 100,000	16.8	9.6	12.5		2018
	95% CI	14.0 - 19.6	7.8 - 11.3	11.0 - 14.1		

Rates are age-adjusted to the U.S. 2000 Standard Population. 95% CI: 95% Confidence Interval of the rate.

Bar Chart shows male and female rates in order listed on table from left to right; dark blue bar indicates the total Maine rate.

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Created March 2020 by Carolyn Bancroft based on work by Mary Caron, Finn Teach, and Sara Huston, Maine Center for Disease Control and Prevention.

The Department of Health and Human Services complies with applicable Federal and State civil rights laws and does not discriminate on the basis of disability, race, color, sex, gender, sexual orientation, age, national origin, religious or political belief, ancestry, familial or marital status, genetic information, association, previous assertion of a claim or right, or whistleblower activity, in admission or access to, or the operation of its policies, programs, services, or activities, or in hiring or employment practices.

Trend At-a-Glance

Leading Causes of Death, Maine, 2009-2018



Indicator	Unit	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	10-year trend	Progress 1999-2018
All Deaths	Rate per 100,000	752.6	749.6	752.8	730.4	754.2	739.0	783.5	759	771.6	753.9		
	95% CI	739.2 - 765.9	736.4 - 762.8	739.6 - 766.0	717.5 - 743.3	741.2 - 767.2	726.2 - 751.8	770.4 - 796.6	746.1 - 772.0	758.7 - 784.5	741.6 - 766.1		
Cancer Deaths	Rate per 100,000	185.2	187.9	182.2	179.3	175.2	170.3	178.0	168.9	170.8	162		
	95% CI	178.7 - 191.8	181.3 - 194.4	175.8 - 188.6	173.1 - 185.6	169.0 - 181.3	164.3 - 176.3	171.9 - 184.1	163 - 174.9	164.9 - 176.7	156.3 - 167.6		
Heart Disease Deaths	Rate per 100,000	156.4	151.1	149.5	145.9	152.3	147.9	157	149.5	143.5	146.9		
	95% CI	150.4 - 162.3	145.3 - 156.9	143.7 - 155.2	140.2 - 151.6	146.5 - 158.0	142.3 - 153.5	151.6 - 163.0	144 - 155.1	138.1 - 148.9	141.6 - 152.3		
Unintentional Injury Deaths	Rate per 100,000	40.9	36.5	38.3	40.4	42.6	45.8	53.8	62.4	68	64.7		
	95% CI	37.6 - 44.3	33.4 - 39.7	35.1 - 41.6	37.0 - 43.7	39.2 - 46.1	42.2 - 49.4	49.9 - 57.7	58.1 - 66.6	63.6 - 72.5	60.4 - 68.9		
Chronic Lower Respiratory Disease Deaths	Rate per 100,000	48.6	47.4	48.4	45.4	49.1	48.1	53.6	47.4	48.7	45.6		
	95% CI	45.3 - 52.0	44.1 - 50.7	45.1 - 51.7	42.2 - 48.5	45.8 - 52.3	44.9 - 51.3	50.3 - 56.9	44.3 - 50.5	45.6 - 52.8	42.6 - 48.5		
Cerebrovascular Disease Deaths	Rate per 100,000	38.0	34.5	35.8	34.3	33.4	33.2	32.6	34.4	37.5	33.8		
	95% CI	35.1 - 41.0	31.7 - 37.3	33.0 - 38.6	31.5 - 37.0	30.7 - 36.0	30.6 - 35.9	30.0 - 35.2	31.7 - 37.1	34.7 - 40.2	31.2 - 36.3		
Alzheimer's Disease Deaths	Rate per 100,000	26.9	28.5	26.8	23.5	21.6	22.7	28.4	29.6	30.4	28.4		
	95% CI	24.5 - 29.4	26.0 - 31.1	24.4 - 29.2	21.2 - 25.7	19.4 - 23.7	20.5 - 24.8	26.0 - 30.8	27.2 - 32.1	27.9 - 32.8	26.0 - 30.7		
Diabetes Deaths	Rate per 100,000	20.6	21.8	20.0	21.3	20.4	22.4	21.9	23.9	19.7	23.4		
	95% CI	18.5 - 22.8	19.6 - 24.1	17.9 - 22.1	19.1 - 23.5	18.3 - 22.5	20.2 - 24.6	19.7 - 24.1	21.7 - 26.2	17.7 - 21.7	21.2 - 25.5		
Influenza and Pneumonia Deaths	Rate per 100,000	14.6	13.4	16.8	11.6	14.0	13.7	17.6	12	15.2	15.2		
	95% CI	12.8 - 16.4	11.7 - 15.1	14.9 - 18.7	10.0 - 13.2	12.3 - 15.8	12.0 - 15.4	15.7 - 19.6	10.4 - 13.6	13.4 - 16.9	13.5 - 16.9		
Suicide Deaths	Rate per 100,000	14.0	13.2	16.6	14.5	17.4	15.7	16.0	15.9	18.9	18.4		
	95% CI	12.0 - 16.0	11.2 - 15.1	14.4 - 18.8	12.5 - 16.6	15.1 - 19.7	13.5 - 17.8	13.9 - 18.2	13.7 - 18.1	16.5 - 21.2	16.1 - 20.7		
Nephritis, Nephrotic Syndrome and Nephrosis Deaths	Rate per 100,000	16.3	15.8	14.8	14.3	13.6	11.7	11.4	12.4	13	12.5		
	95% CI	14.3 - 18.2	13.9 - 17.7	12.9 - 16.6	12.5 - 16.1	11.9 - 15.2	10.1 - 13.3	9.8 - 13.0	10.8 - 14	11.4 - 14.5	11.0 - 14.1		

■ = Improving

● = Plateaued

◆ = Worsening

Rates are age-adjusted to the U.S. 2000 Standard Population.

95% CI: 95% Confidence Interval of the rate or percent.

"Progress" column is based on Joinpoint regression analysis of 1999-2018 WONDER data showing statistically significant trend (improving or worsening), or no significant trend (plateaued) for most recent joinpoint segment.

"Trend" column shows trend line for period shown; dark marker indicates the best rate.

Data Source: United States Department of Health and Human Services (US DHHS), Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death 1999-2018 on CDC WONDER Online Database, released December 2018. Data are from the Multiple Cause of Death Files, 1999- 2018 as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Created February 2020 by Carolyn Bancroft, Maine Center for Disease Control and Prevention building on work by Mary Caron, Finn Teach, and Sara Huston.

The Department of Health and Human Services complies with applicable Federal and State civil rights laws and does not discriminate on the basis of disability, race, color, sex, gender, sexual orientation, age, national origin, religious or political belief, ancestry, familial or marital status, genetic information, association, previous assertion of a claim or right, or whistleblower activity, in admission or access to, or the operation of its policies, programs, services, or activities, or in hiring or employment practices.

County At-a-Glance

Leading Causes of Death, Maine, 2014-2018, 5-year average



Indicator	Unit	Androscoggin	Aroostook	Cumberland	Franklin	Hancock	Kennebec	Knox	Lincoln	Oxford	Penobscot	Piscataquis	Sagadahoc	Somerset	Waldo	Washington	York	Maine Total	Bar Chart
All Deaths	Rate per 100,000	837.5	793.6	687.4	806.2	714.8	808.1	731.2	711.7	777.2	808.5	919.3	717.3	868.3	744.9	896.7	721.4	760.9	
	95% CI	816.1 - 859.0	769.7 - 818.1	675.8 - 699.0	768.1 - 844.3	688.9 - 740.8	788.8 - 827.5	701.4 - 761.0	679.6 - 743.7	750.4 - 804.0	791.0 - 825.9	866.6 - 972.0	684.6 - 750.0	838.3 - 900.3	712.2 - 777.5	837.7 - 899.6	707.4 - 735.5	755.3 - 766.5	
Cancer Deaths	Rate per 100,000	178.4	166.8	153.5	181.0	157.8	176.5	175.0	156.4	179.4	171.0	215.6	170.2	196.7	165.5	197.6	169.5	169.6	
	95% CI	168.4 - 188.4	155.6 - 178.0	147.9 - 159.1	163.0 - 199.1	145.8 - 169.8	167.5 - 185.5	160.4 - 189.5	142.0 - 170.8	166.5 - 192.3	162.9 - 179.1	191.8 - 239.5	154.4 - 185.9	182.1 - 211.3	150.3 - 180.8	179.8 - 215.3	162.7 - 176.3	167.0 - 172.2	
Heart Disease Deaths	Rate per 100,000	166.0	160.0	124.8	152.8	142.8	158.7	153.3	147.2	138.5	161.1	190.4	147.9	195.1	169.9	179.3	131.6	148.5	
	95% CI	156.4 - 175.6	149.5 - 170.5	119.9 - 129.8	136.3 - 169.2	131.4 - 154.2	150.2 - 167.3	140.4 - 166.2	133.2 - 161.2	127.2 - 149.8	153.3 - 168.9	167.3 - 213.5	133.3 - 162.5	180.5 - 209.7	154.4 - 185.4	162.5 - 196.2	125.7 - 137.5	146.1 - 151.0	
Unintentional Injury Deaths	Rate per 100,000	56.9	47.3	52.8	55.4	59.0	62.2	60.3	58.4	51.9	60.0	82.5	41.0	66.7	59.4	98.6	65.6	58.8	
	95% CI	50.6 - 63.3	40.0 - 54.7	49.2 - 56.5	43.9 - 68.9	49.8 - 68.2	55.9 - 68.5	49.7 - 70.8	47.0 - 71.7	43.5 - 60.4	54.5 - 65.6	63.3 - 105.8	32.5 - 51.1	56.5 - 76.9	48.4 - 70.3	82.0 - 115.1	60.6 - 70.6	57.0 - 60.7	
Chronic Lower Respiratory Disease Deaths	Rate per 100,000	57.4	59.2	38.4	55.1	41.8	51.7	44.5	35.5	65.9	56.1	56.3	40.0	69.0	42.0	53.3	42.2	48.5	
	95% CI	51.7 - 63.1	52.9 - 65.6	35.6 - 4.1	45.4 - 64.8	35.6 - 47.9	46.9 - 56.6	37.1 - 51.9	29.0 - 42.0	58.2 - 73.6	51.5 - 60.8	44.9 - 69.8	32.5 - 47.6	60.3 - 77.6	34.5 - 49.5	44.1 - 62.5	38.9 - 45.5	47.1 - 49.9	
Cerebrovascular Disease Deaths	Rate per 100,000	38.7	34.9	31.8	39.9	34.3	40.9	30.4	42.3	28.1	34.5	40.9	45.7	33.8	37.4	32.3	27.7	34.2	
	95% CI	34.0 - 43.4	30.0 - 39.7	29.3 - 34.3	32.1 - 49.1	28.8 - 39.8	36.5 - 45.2	24.6 - 36.3	35.2 - 49.5	23.0 - 33.1	30.9 - 38.1	30.7 - 53.6	37.4 - 54.0	27.7 - 40.0	30.4 - 45.6	25.5 - 40.4	25.0 - 30.4	33.0 - 35.3	
Alzheimer's Disease Deaths	Rate per 100,000	54.5	24.3	27.5	24.3	24.5	25.9	27.9	19.3	27.8	20.3	29.6	13.7	22.3	22.5	19.7	33.5	27.9	
	95% CI	49.0 - 60.0	20.4 - 28.2	25.2 - 29.8	18.2 - 31.8	19.9 - 29.0	22.5 - 29.4	22.5 - 33.3	14.9 - 24.6	22.8 - 32.8	17.5 - 23.0	21.5 - 39.7	9.6 - 18.6	17.6 - 27.9	17.2 - 29.0	14.6 - 25.9	30.6 - 36.5	26.8 - 28.9	
Diabetes Deaths	Rate per 100,000	24.2	25.2	16.3	34.0	21.1	27.3	18.7	20.9	35.1	24.2	25.5	12.6	24.9	20.2	19.8	21.1	22.2	
	95% CI	20.4 - 27.9	21.0 - 29.3	14.5 - 18.1	26.2 - 43.4	16.8 - 26.0	23.7 - 31.0	14.3 - 24.0	15.6 - 27.4	29.4 - 40.9	21.2 - 27.2	18.0 - 35.2	8.5 - 18.0	20.0 - 30.6	15.2 - 26.3	14.9 - 25.8	18.7 - 23.5	21.3 - 23.2	
Influenza and Pneumonia Deaths	Rate per 100,000	17.3	17.0	13.4	19.1	14.0	14.1	10.6	11.6	14.1	17.9	14.1	14.6	13.8	14.9	21.7	12.9	14.7	
	95% CI	14.2 - 20.4	13.8 - 20.8	11.8 - 15.1	13.3 - 26.6	10.6 - 18.1	11.2 - 16.6	7.2 - 14.9	8.1 - 16.1	10.8 - 18.2	15.3 - 20.4	8.2 - 22.6	10.5 - 19.9	10.0 - 18.4	10.6 - 20.3	16.3 - 28.2	11.1 - 14.8	13.9 - 15.5	
Suicide Deaths	Rate per 100,000	17.8	18.7	12.3	19.0	15.8	19.3	17.9	20.2	15.2	16.9	23.2	23.2	20.9	19.6	26.0	16.8	16.9	
	95% CI	14.4 - 21.7	14.2 - 24.2	10.5 - 14.1	12.5 - 27.6	11.4 - 21.2	15.8 - 22.9	12.3 - 25.1	13.6 - 28.8	11.0 - 20.4	13.9 - 19.8	13.5 - 37.1	16.4 - 31.8	15.6 - 27.4	13.9 - 26.8	18.2 - 36.0	14.2 - 19.4	15.9 - 17.9	
Nephritis, Nephrotic Syndrome and Nephrosis Deaths	Rate per 100,000	8.7	15.3	12.1	14.9	12.8	11.1	9.8	7.6	12.9	16.2	8.4	13.7	8.1	10.6	14.1	12.3	12.2	
	95% CI	6.6 - 11.2	12.3 - 18.9	10.5 - 13.6	10.3 - 21.0	9.7 - 16.5	9.0 - 13.6	6.8 - 13.6	4.6 - 11.7	9.8 - 16.7	13.7 - 18.6	4.3 - 14.7	9.7 - 18.9	5.4 - 11.6	6.9 - 15.7	9.6 - 19.9	10.5 - 14.2	10.5 - 14.2	

Orange shading indicates county rate is significantly worse than the state rate.
 Green shading indicates county rate is significantly better than the state rate.

DNR: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

Rates are age-adjusted to the U.S. 2000 Standard Population.

95% CI: 95% Confidence Interval of the rate.

Bar Chart shows county rates in order listed on table from left to right; dark blue bar is the Maine Total rate.

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. March 2020

Created November 2019 by Carolyn Bancroft based on work by Mary Caron, Finn Teach, and Sara Huston, Maine Center for Disease Control and Prevention.

The Department of Health and Human Services complies with applicable Federal and State civil rights laws and does not discriminate on the basis of disability, race, color, sex, gender, sexual orientation, age, national origin, religious or political belief, ancestry, familial or marital status, genetic information, association, previous assertion of a claim or right, or whistleblower activity, in admission or access to, or the operation of its policies, programs, services, or activities, or in hiring or employment practices.

Public Health Districts, At-a-Glance

Leading Causes of Death, Maine, 2014-2018



Indicator	Unit	Maine Public Health Districts									Bar Chart	Years
		Aroostook	Downeast	Penquis	Central Maine	Mid Coast Maine	Western Maine	Cumberland	York	Maine Total		
All Deaths	Rate per 100,000	793.9	780.9	820.3	826.7	725.1	813.4	687.4	721.4	761.3		2014-2018
	95% CI	769.7 - 818.1	759.2 - 802.6	803.6 - 836.8	810.3 - 843.1	709.4 - 740.9	798.1 - 828.7	675.8 - 699.0	707.4 - 735.5	755.7 - 766.9		
Cancer Deaths	Rate per 100,000	166.8	172.0	176.8	182.5	166.6	179.0	153.5	169.5	169.5		2014-2018
	95% CI	155.6 - 178.0	162.0 - 182.0	169.1 - 184.4	174.8 - 190.1	159.2 - 174.1	171.8 - 186.2	147.9 - 159.1	162.7 - 176.3	166.9 - 172.2		
Heart Disease Deaths	Rate per 100,000	160.0	156.0	164.6	169.5	154.1	155.1	124.8	131.6	148.6		2014-2018
	95% CI	149.5 - 170.5	146.5 - 165.5	157.2 - 172.0	162.0 - 176.9	147.1 - 161.2	148.4 - 161.8	119.9 - 129.8	125.7 - 137.5	146.1 - 151.0		
Unintentional Injury Deaths	Rate per 100,000	47.3	73.1	61.9	63.5	54.6	55.2	52.8	65.6	58.9		2014-2018
	95% CI	40.0 - 54.7	64.7 - 81.5	56.6 - 67.3	58.1 - 68.9	49.3 - 59.9	50.5 - 59.9	49.2 - 56.5	60.6 - 70.6	57.0 - 60.7		
Chronic Lower Respiratory Disease Deaths	Rate per 100,000	59.2	46.0	56.2	57.0	40.7	59.6	38.4	42.2	48.5		2014-2018
	95% CI	51.8 - 60.5	40.8 - 51.1	51.8 - 60.5	52.7 - 61.2	37.1 - 44.3	55.5 - 63.8	35.6 - 41.1	38.9 - 45.5	47.1 - 49.9		
Cerebrovascular Disease Deaths	Rate per 100,000	34.9	33.8	35.2	38.8	38.8	35.4	31.8	27.7	34.2		2014-2018
	95% CI	20.0 - 39.7	29.4 - 38.2	31.8 - 38.7	35.2 - 42.4	35.2 - 42.3	32.2 - 38.6	29.3 - 34.3	25.0 - 30.4	33.0 - 35.3		
Alzheimer's Disease Deaths	Rate per 100,000	24.3	22.8	21.4	24.9	21.4	41.1	27.5	33.5	27.9		2014-2018
	95% CI	20.4 - 28.2	19.2 - 26.3	18.8 - 24.0	22.1 - 27.7	18.8 - 24.0	37.6 - 44.5	25.2 - 29.8	30.6 - 36.5	26.8 - 28.9		
Diabetes Deaths	Rate per 100,000	25.2	20.6	24.3	26.6	18.3	29.3	16.3	21.1	22.2		2014-2018
	95% CI	21.0 - 29.3	17.2 - 24.1	21.5 - 27.2	23.6 - 29.6	15.7 - 20.8	26.3 - 32.3	14.5 - 18.1	18.7 - 23.5	21.3 - 23.2		
Influenza and Pneumonia Deaths	Rate per 100,000	17.0	16.8	17.3	14.1	12.5	16.5	13.4	12.9	14.7		2014-2018
	95% CI	13.8 - 20.8	13.7 - 19.9	14.9 - 19.7	11.9 - 16.2	10.5 - 14.5	14.3 - 18.8	11.8 - 15.1	11.1 - 14.8	13.9 - 15.5		
Suicide Deaths	Rate per 100,000	18.7	19.5	17.4	19.9	20.0	17.3	12.3	16.8	16.9		2014-2018
	95% CI	14.2 - 24.2	15.4 - 24.3	14.6 - 20.3	16.8 - 22.9	16.7 - 23.4	14.7 - 19.9	10.5 - 14.1	14.2 - 19.4	15.9 - 17.9		
Nephritis, Nephrotic Syndrome and Nephrosis Deaths	Rate per 100,000	15.3	13.3	15.2	10.2	10.3	11.1	12.1	12.3	12.2		2014-2018
	95% CI	12.3 - 18.9	10.7 - 16.4	13.0 - 17.5	8.4 - 12.0	8.4 - 12.2	9.3 - 12.8	10.5 - 13.6	10.5 - 14.2	11.5 - 12.9		

Orange shading indicates county rate is significantly worse than the state rate.
 Green shading indicates county rate is significantly better than the state rate.

Rates are age-adjusted to the U.S. 2000 Standard Population.

95% CI: 95% Confidence Interval of the rate.

Bar Chart shows county rates in order listed on table from left to right; dark blue bar is the Maine Total rate.

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. December 2019.

Created November 2019 by Carolyn Bancroft based on work by Mary Caron, Finn Teach, and Sara Huston, Maine Center for Disease Control and Prevention.

The Department of Health and Human Services complies with applicable Federal and State civil rights laws and does not discriminate on the basis of disability, race, color, sex, gender, sexual orientation, age, national origin, religious or political belief, ancestry, familial or marital status, genetic information, association, previous assertion of a claim or right, or whistleblower activity, in admission or access to, or the operation of its policies, programs, services, or activities, or in hiring or employment practices.

Age Groups At-a-Glance

Leading Causes of Death, Maine, 2018



Indicator	Unit	Under 5 Yrs	5 - 14 Yrs	15 - 24 Yrs	25 - 34 Yrs	35 - 44 Yrs	45 - 54 Yrs	55 - 64 Yrs	65 - 74 Yrs	75 - 84 Yrs	85+ Yrs	Bar Chart	Year
All Deaths	Rate per 100,000	107.3	13.5	63.0	139.1	212.6	399.6	823.7	1702.9	4,750.5	14,880.8		2018
	95% CI	82.0-132.7	7.4-19.6	50.4-75.6	120.8-157.3	189.5-235.8	370.4-428.8	785.0-862.3	1639.6-1766.2	4598.7-4902.3	14467.2-15294.4		
Cancer Deaths	Rate per 100,000	*	*	*	DNR	28.2	112.7	288.7	560.1	1,151.3	1,684.7		2018
	95% CI	*	*	*	3.7-12.5	19.8-36.7	97.2-128.2	265.8-311.6	523.8-596.4	1076.6-1226.1	1545.5-1823.8		
Heart Disease Deaths	Rate per 100,000	*	*	*	DNR	28.2	62.2	134.4	304.2	924.1	3,824.2		2018
	95% CI	*	*	*	2.8 - 10.9	19.8-36.7	50.6 - 73.7	118.8 - 150.1	277.5 - 331.0	857.1 - 991.0	3614.5 - 4033.8		
Unintentional Injury Deaths	Rate per 100,000	*	*	25.6	74.2	82.7	63.8	54.7	52.6	175.5	715.2		2018
	95% CI	*	*	17.6 - 33.6	60.9 - 87.6	68.2 - 97.1	52.1 - 75.5	44.8 - 64.7	41.5 - 63.8	146.3 - 204.6	624.5 - 805.8		
Chronic Lower Respiratory Disease Deaths	Rate per 100,000	*	*	*	*	*	DNR	56.6	134.7	392.6	787.0		2018
	95% CI	*	*	*	*	*	5.8-15.3	46.5-66.7	116.9-152.5	349.0-436.2	691.9-882.1		
Cerebrovascular Disease Deaths	Rate per 100,000	*	*	*	*	*	DNR	27.8	61.8	229.8	984.5		2018
	95% CI	*	*	*	*	*	3.3-11.1	20.7-34.9	49.8-73.9	196.4-263.1	878.1-1090.9		
Alzheimer's Disease Deaths	Rate per 100,000	*	*	*	*	*	*	*	25.7	209.6	1,095.2		2018
	95% CI	*	*	*	*	*	*	*	18.0-33.5	177.7-241.4	983.0-1207.4		
Diabetes Deaths	Rate per 100,000	*	*	*	*	*	16.1	40.1	82.0	165.4	284.3		2018
	95% CI	*	*	*	*	*	10.2 - 22.0	31.6 - 48.6	68.1-95.9	137.1-193.7	227.1-341.4		
Influenza and Pneumonia Deaths	Rate per 100,000	*	*	*	*	*	*	14.6	33.7	88.4	424.9		2018
	95% CI	*	*	*	*	*	*	9.5-19.8	24.8-42.6	67.7-109.1	355.0-494.8		
Suicide Deaths	Rate per 100,000	*	*	19.0	22.5	22.3	26.6	26.4	21.4	26.5	*		2018
	95% CI	*	*	12.1-26.0	15.1-29.8	14.8-29.8	19.1-34.2	19.5-33.3	14.3-28.5	15.2-37.8	*		
Nephritis, Nephrotic Syndrome and Nephrosis Deaths	Rate per 100,000	*	*	*	*	*	*	12.7	29.4	84.6	323.2		2018
	95% CI	*	*	*	*	*	*	7.9-17.5	21.1-37.7	64.3-104.8	262.2-384.1		

Rates are crude rates, deaths per 100,000 population.

DNR: Data not reliable; Using the WONDER system definition, death rates are noted as unreliable when the numerator is 20 or less. * Counts less than 10 are suppressed

95% CI: 95% Confidence Interval of the rate.

Bar Chart shows age group rates in order listed on table from left to right.

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. March 2020.







Created December 202 by Anne Rogers and Carolyn Bancroft based on work by Mary Caron, Finn Teach, and Sara Huston, Maine Center for Disease Control and Prevention.

The Department of Health and Human Services complies with applicable Federal and State civil rights laws and does not discriminate on the basis of disability, race, color, sex, gender, sexual orientation, age, national origin, religious or political belief, ancestry, familial or marital status, genetic information, association, previous assertion of a claim or right, or whistleblower activity, in admission or access to, or the operation of its policies, programs, services, or activities, or in hiring or employment practices.

Section 3:

Cause-specific Figures

Death Rate Trends, Maine, 1999-2018








Data Sources: CDC WONDER (1999-2017 data). 2018 data: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. October 2019- January 2020.

Rates are deaths per 100,000 population, age-adjusted to the U.S. 2000 standard population.

See Section 5: Tables for detailed data.

Death Rate Trends, Maine, 1999-2018









Data Sources: CDC WONDER (1999-2017 data). 2018 data: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. October 2019- January 2020.

Rates are deaths per 100,000 population, age-adjusted to the U.S. 2000 standard population.

See Section 5: Tables for detailed data.

Death Rate Trends, Maine, 1999-2018








Data Sources: CDC WONDER (1999-2017 data). 2018 data: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. January 2020

Rates are deaths per 100,000 population, age-adjusted to the U.S. 2000 standard population.

See Section 5: Tables for detailed data.

Death Rate Trends, Maine, 1999-2018







Data Sources: CDC WONDER (1999-2017 data). 2018 data: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. January 2020.

Rates are deaths per 100,000 population, age-adjusted to the U.S. 2000 standard population.

See Section 5: Tables for detailed data.

Death Rate Trends by Race and Ethnicity, Maine, 1999-2018

Please note the different scale in these figures.



Data Source: CDC WONDER (1999-2018 data).


Rates are deaths per 100,000 population, age-adjusted to the U.S. 2000 standard population. See Section 5: Tables for detailed data.

Section 4:


Cause-specific Maps

Death Rates by Cause and County of Residence, Maine, 2014-2018


All Causes




Cancer




Heart Disease




Cerebrovascular Disease



Unintentional Injury




Chronic Lower Respiratory Disease



Quartile based on county rate:


- 1st Quartile (Lowest)
- 2nd Quartile
- 3rd Quartile
- 4th Quartile (Highest)




Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. Rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population. Death rates mapped using the quantiles method with four categories. Map created by Finn Teach, February, 2020. See Section 5: Tables for detailed data.

Death Rates by Cause and County of Residence, Maine, 2014-2018


Diabetes




Influenza & Pneumonia




Alzheimer's Disease



Suicide




Nephritis, Nephrotic Syndrome, and Nephrosis



Quartile based on county rate:

- 1st Quartile (Lowest)
- 2nd Quartile
- 3rd Quartile
- 4th Quartile (Highest)



Data Source: Maine Center for Disease Control and Prevention. Office of Data, Research and Vital Statistics. Rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population. Death rates mapped using the quantiles method with four categories. Map created by Finn Teach, February, 2020. See Section 5: Tables for detailed data.

Section 5:

Detailed Tables

List of Tables

Table	Title
1	<u>Leading Causes of Death in Maine, with U.S. Comparison Rates, 2018</u>
2	<u>Leading Causes of Death by Sex, Maine, 2018</u>
3	<u>Number of Deaths due to Leading Causes of Death by Age Group, Maine, 2018</u>
4	<u>Leading Causes of Death by Race, Maine, 2009-2018</u>
5	<u>Leading Causes of Death by Hispanic Ethnicity, Maine, 2009-2018</u>
6	<u>Leading Causes of Death by County, Maine, 2018</u>
7	<u>Years of Potential Life Lost (YPLL) before Age 75, by Cause of Death, Maine vs. U.S, 2018</u>
8	<u>All Deaths by Year, Maine and U.S, 1999-2018</u>
9	<u>All Deaths by Sex and Year, Maine, 1999-2018</u>
10	<u>All Deaths by Sex and Age, Maine, 2018</u>
11	<u>All Deaths by Race and Hispanic Ethnicity, Maine, 2009-2018</u>
12	<u>All Deaths by County, Maine, 2014-2018</u>
13	<u>Cancer Deaths by Year, Maine and U.S, 1999-2018</u>
14	<u>Cancer Deaths by Sex and Year, Maine, 1999-2018</u>
15	<u>Cancer Deaths by Sex and Age, Maine, 2018</u>
16	<u>Cancer Deaths by Race and Hispanic Ethnicity, Maine, 2009-2018</u>
17	<u>Cancer Deaths by County, Maine, 2014-2018</u>
18	<u>Heart Disease Deaths by Year, Maine and U.S, 1999-2018</u>
19	<u>Heart Disease Deaths by Sex and Year, Maine, 1999-2018</u>
20	<u>Heart Disease Deaths by Sex and Age, Maine, 2018</u>
21	<u>Heart Disease Deaths by Race and Hispanic Ethnicity, Maine, 2009-2018</u>
22	<u>Heart Disease Deaths by County, Maine, 2014-2018</u>
23	<u>Chronic Lower Respiratory Disease Deaths by Year, Maine and U.S, 1999-2018</u>
24	<u>Chronic Lower Respiratory Disease Deaths by Sex and Year, Maine, 1999-2018</u>
25	<u>Chronic Lower Respiratory Disease Deaths by Sex and Age, Maine, 2018</u>
26	<u>Chronic Lower Respiratory Disease Deaths by Race and Hispanic Ethnicity, Maine, 2009-2018</u>
27	<u>Chronic Lower Respiratory Disease Deaths by County, Maine, 2014-2018</u>
28	<u>Unintentional Injury Deaths by Year, Maine and U.S, 1999-2018</u>
29	<u>Unintentional Injury Deaths by Sex and Year, Maine, 1999-2018</u>
30	<u>Unintentional Injury Deaths by Sex and Age, Maine, 2018</u>
31	<u>Unintentional Injury Deaths by Race and Hispanic Ethnicity, Maine, 2009-2018</u>
32	<u>Unintentional Injury Deaths by County, Maine, 2014-2018</u>
33	<u>Cerebrovascular Disease Deaths by Year, Maine and U.S, 1999-2018</u>
34	<u>Cerebrovascular Disease Deaths by Sex and Year, Maine, 1999-2018</u>
35	<u>Cerebrovascular Disease Deaths by Sex and Age, Maine, 2018</u>
36	<u>Cerebrovascular Disease Deaths by Race and Hispanic Ethnicity, Maine, 2009-2018</u>
37	<u>Cerebrovascular Disease Deaths by County, Maine, 2014-2018</u>

- 38 [Alzheimer's Disease Deaths by Year, Maine and U.S, 1999-2018](#)
- 39 [Alzheimer's Disease Deaths by Sex and Year, Maine, 1999-2018](#)
- 40 [Alzheimer's Disease Deaths by Sex and Age, Maine, 2018](#)
- 41 [Alzheimer's Disease Deaths by Race and Hispanic Ethnicity, Maine, 2009-2018](#)
- 42 [Alzheimer's Disease Deaths by County, Maine, 2014-2018](#)
- 43 [Diabetes Deaths by Year, Maine and U.S, 1999-2018](#)
- 44 [Diabetes Deaths by Sex and Year, Maine, 1999-2018](#)
- 45 [Diabetes Deaths by Sex and Age, Maine, 2018](#)
- 46 [Diabetes Deaths by Race and Hispanic Ethnicity, Maine, 2009-2018](#)
- 47 [Diabetes Deaths by County, Maine, 2014-2018](#)
- 48 [Influenza and Pneumonia Disease Deaths by Year, Maine and U.S, 1999-2018](#)
- 49 [Influenza and Pneumonia Deaths by Sex and Year, Maine, 1999-2018](#)
- 50 [Influenza and Pneumonia Deaths by Sex and Age, Maine, 2018](#)
- 51 [Influenza and Pneumonia Deaths by Race and Hispanic Ethnicity, Maine, 2009-2018](#)
- 52 [Influenza and Pneumonia Deaths by County, Maine, 2014-2018](#)
- 53 [Suicide Deaths by Year, Maine and U.S, 1999-2018](#)
- 54 [Suicide Deaths by Sex and Year, Maine, 1999-2018](#)
- 55 [Suicide Deaths by Sex and Age, Maine, 2018](#)
- 56 [Suicide Deaths by Race and Hispanic Ethnicity, Maine, 2009-2018](#)
- 57 [Suicide Deaths by County, Maine, 2014-2018](#)
- 58 [Nephritis, Nephrotic Syndrome, and Nephrosis Deaths by Year, Maine and U.S, 1999-2018](#)
- 59 [Nephritis, Nephrotic Syndrome, and Nephrosis Deaths by Sex and Year, Maine, 1999-2018](#)
- 60 [Nephritis, Nephrotic Syndrome, and Nephrosis Deaths by Sex and Age, Maine, 2018](#)
- 61 [Nephritis, Nephrotic Syndrome, and Nephrosis Deaths by Race and Hispanic Ethnicity, Maine, 2009-2018](#)
- 62 [Nephritis, Nephrotic Syndrome, and Nephrosis Deaths by County, Maine, 2014-2018](#)

Table 1. Leading Causes of Death in Maine, with U.S. Comparison Rates, 2018

Rank	Cause	Number of Deaths	Maine		U.S.		Rank
			Age-adjusted Rate		Age-adjusted Rate		
			Rate	95% CI	Rate	95% CI	
1	Cancer (C00-C97)	3,273	162.0	156.3 - 167.6	149.1	148.7 - 149.5	2
2	Heart disease (I00-I09,I11,I13,I20-I51)	2,963	146.9	141.6 - 152.3	163.6	163.2 - 164.0	1
3	Unintentional injury (V01-X59,Y85-Y86)	996	64.7	60.4 - 68.9	48	47.8 - 48.2	3
4	Chronic lower respiratory disease (J40-J47)	937	45.6	42.6 - 48.5	39.7	39.5 - 39.9	4
5	Cerebrovascular disease (I60-I69)	691	33.8	31.2 - 36.3	37.1	36.9 - 37.3	5
6	Alzheimer's disease (G30)	580	28.4	26.0 - 30.7	30.5	30.3 - 30.7	6
7	Diabetes (E10-E14)	479	23.4	21.3 - 25.6	21.4	21.2 - 21.5	7
8	Influenza and pneumonia (J09-J18)	311	15.2	13.5 - 16.9	14.9	14.8 - 15.0	8
9	Suicide (X60-X84,Y87.0)	269	18.4	16.1 - 20.7	14.2	14.1 - 14.3	10
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	258	12.5	11.0 - 14.1	12.9	12.8 - 13.0	9

Maine Data Source: Maine CDC, Data Research, and Vital Statistics. US Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death. 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Feb 13, 2020

ICD-10 codes are grouped by the National Center for Health Statistics Top 15 Rankable Cause List.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 2. Leading Causes of Death by Sex, Maine, 2018

Males					
Rank	Cause	Number of Deaths	Crude Rate	Age-adjusted Rate	
				Rate	95% CI
1	Cancer (C00-C97)	1,717	262.0	188.8	179.7 - 198.0
2	Heart disease (I00-I09,I11,I13,I20-I51)	1,648	251.5	193.7	184.2 - 203.2
3	Unintentional injury (V01-X59,Y85-Y86)	615	93.7	89.7	82.3 - 97.1
4	Chronic lower respiratory disease (J40-J47)	476	72.6	53.5	48.6 - 58.4
5	Cerebrovascular disease (I60-I69)	275	42.0	32.0	28.1 - 35.8
6	Diabetes (E10-E14)	271	41.4	29.4	25.8 - 33.0
7	Suicide (X60-X84,Y87.0)	217	33.1	30.4	26.1 - 34.6
8	Influenza and pneumonia (J10-J18)	168	25.6	20.1	16.9 - 23.2
9	Alzheimer's disease (G30)	164	25.0	20.4	17.3 - 23.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	21.8	16.8	14.0 - 19.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	132	20.1	14.9	12.3 - 17.6
12	Parkinson's disease (G20-G21)	115	17.5	13.9	11.4 - 16.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	58	8.8	6.9	5.2 - 8.9
14	Septicemia (A40-A41)	53	8.1	6.1	4.5 - 8.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	7.9	6.2	4.6 - 8.2
Females					
Rank	Cause	Number of Deaths	Crude Rate	Age-adjusted Rate	
				Rate	95% CI
1	Cancer (C00-C97)	1,556	227.8	141.1	133.9 - 148.4
2	Heart disease (I00-I09,I11,I13,I20-I51)	1,315	192.5	110.0	103.9 - 116.1
3	Chronic lower respiratory disease (J40-J47)	462	67.6	39.7	36.0 - 43.3
4	Cerebrovascular disease (I60-I69)	416	60.9	34.1	30.7 - 37.4
5	Alzheimer's disease (G30)	416	60.9	33.2	29.9 - 36.4
6	Unintentional injury (V01-X59,Y85-Y86)	381	55.8	40.7	36.3 - 45.2
7	Diabetes (E10-E14)	208	30.5	18.1	15.6 - 20.7
8	Influenza and pneumonia (J10-J18)	143	20.9	11.7	9.8 - 13.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	16.8	9.6	7.8 - 11.3
10	Parkinson's disease (G20-G21)	72	10.5	6.1	4.7 - 7.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	66	9.7	6.9	5.2 - 8.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	65	9.5	5.1	3.9 - 6.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	7.8	4.6	3.4 - 6.2
14	Suicide (X60-X84,Y87.0)	52	7.6	7.1	5.2 - 9.5
15	Septicemia (A40-A41)	38	5.6	6.4	2.3 - 4.7

Data Source: Maine CDC, Data Research, and Vital Statistics.

ICD-10 codes are grouped by ICD-10 113 Cause List.

Crude rates are deaths per 100,000 population.

95% CI: 95% confidence interval of the rate.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 3. Number of Deaths due to Leading Causes of Death, by Age Group, Maine, 2018

Age Groups	Rank									
	1	2	3	4	5	6	7	8	9	10
<1	Congenital Anomalies 13	Unintentional Injury 11	Maternal Pregnancy Comp. ---	Unintentional Injury ---	Placenta Cord Membranes ---	Circulatory System Disease ---	Respiratory Distress ---	SIDS ---	Six Tied ---	Six Tied ---
1-4	Congenital Anomalies ---	Pneumonitis ---								
5-9	Unintentional Injury ---	Congenital Anomalies ---	Suicide ---							
10-14	Unintentional Injury ---	Cerebrovascular Disease ---	Chronic Lower Respiratory Disease ---	Heart Disease ---	Homicide ---	Cancer ---	Suicide ---			
15-24	Unintentional Injury 39	Suicide 29	Cancer ---	Heart Disease ---	Congenital Anomalies ---	Chronic Lower Respiratory Disease ---	Homicide ---	Septicemia ---		
25-34	Unintentional Injury 120	Suicide 37	Cancer 13	Heart Disease 11	Influenza & Pneumonia ---	Cerebrovascular Disease ---	Nephritis ---	Seven Tied ---	Seven Tied ---	Seven Tied ---
35-44	Unintentional Injury 126	Heart Disease 43	Cancer 43	Suicide 34	Liver Disease 12	Diabetes ---	Homicide ---	Influenza & Pneumonia ---	Three Tied ---	Three Tied ---
45-54	Cancer 204	Unintentional Injury 117	Heart Disease 113	Suicide 48	Liver Disease 34	Diabetes 29	Chronic Lower Respiratory Disease 19	Cerebrovascular Disease 13	Septicemia ---	Three Tied ---
55-64	Cancer 611	Heart Disease 286	Chronic Lower Respiratory Disease 120	Unintentional Injury 116	Diabetes 85	Cerebrovascular Disease 59	Suicide 56	Liver Disease 53	Influenza & Pneumonia 31	Nephritis 27
65+	Heart Disease 2,507	Cancer 2,392	Chronic Lower Respiratory Disease 794	Cerebrovascular Disease 611	Alzheimer's Disease 574	Unintentional Injury 465	Diabetes 359	Influenza & Pneumonia 268	Nephritis 223	Parkinson's Disease 187
All Ages	Cancer 3,275	Heart Disease 2,965	Unintentional Injury 1,000	Chronic Lower Respiratory Disease 938	Cerebrovascular Disease 690	Alzheimer's Disease 580	Diabetes 478	Influenza & Pneumonia 312	Suicide 270	Nephritis 258

Note: For leading cause categories in this State-level chart, counts of less than 10 deaths have been suppressed (---). "Tied" refers to a tie among multiple causes.
 Data Source: National Center for Health Statistics (NCHS), National Vital Statistics System. Centers for Disease Control and Prevention: WISQARS Leading Causes of Death. Available at: www.cdc.gov/injury/wisqars

Table 4. Leading Causes of Death by Race, Maine, 2009-2018 (page 1 of 2)

Cause of Death	Total Number of Deaths	Crude Rate	Age-adjusted Rate	
			Rate	95% CI
American Indian or Alaska Native				
1 Cancer (C00-C97)	160	147.1	190.0	156.9 - 223.2
2 Heart disease (I00-I09,I11,I13,I20-I51)	104	95.6	163.1	127.5 - 198.6
3 Unintentional injury (V01-X59,Y85-Y86)	71	65.3	70.9	54.1 - 91.3
4 Chronic lower respiratory disease (J40-J47)	67	61.6	85.0	64.0 - 110.6
5 Diabetes (E10-E14)	38	34.9	50.7	34.0 - 72.9
6 Suicide (X60-X84,Y87.0)	21	19.3	22.8	13.3 - 36.5
7 Cerebrovascular disease (I60-I69)	20	18.4	37.6	21.9 - 60.1
8 Chronic liver disease and cirrhosis (K70,K73-K74)	14	DNR	DNR	6.9 - 21.2
9 #Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	DNR	DNR	6.7 - 34.5
Asian or Pacific Islander				
1 Cancer (C00-C97)	100	58.8	84.6	66.9 - 102.2
2 Heart disease (I00-I09,I11,I13,I20-I51)	59	34.7	62.1	46.2 - 81.6
3 Cerebrovascular disease (I60-I69)	32	18.8	34.2	23.1 - 48.8
4 Unintentional injury (V01-X59,Y85-Y86)	27	15.9	21.8	13.7 - 33.1
5 Diabetes (E10-E14)	15	DNR	DNR	7.7 - 24.6
6 Alzheimer's disease (G30)	12	DNR	DNR	7.8 - 27.9
7 Suicide (X60-X84,Y87.0)	11	DNR	DNR	3.2 - 11.3
Black or African American				
1 Cancer (C00-C97)	122	53.9	118.0	95.0 - 141.0
2 Heart disease (I00-I09,I11,I13,I20-I51)	93	41.1	97.9	77.4 - 122.2
3 Unintentional injury (V01-X59,Y85-Y86)	59	26.0	33.2	24.1 - 44.5
4 Certain conditions originating in the perinatal period (P00-P96)	23	10.2	6.9	4.4 - 10.4
5 Suicide (X60-X84,Y87.0)	23	10.2	10.0	6.2 - 15.3
6 Diabetes (E10-E14)	20	8.8	17.5	10.0 - 28.3
7 Cerebrovascular disease (I60-I69)	19	DNR	DNR	12.1 - 35.6
8 Chronic lower respiratory disease (J40-J47)	14	DNR	DNR	9.3 - 31.5
9 Human immunodeficiency virus (HIV) disease (B20-B24)	14	DNR	DNR	5.3 - 18.0
10 Homicide (*U01-*U02,X85-Y09,Y87.1)	12	DNR	DNR	2.7 - 9.8
11 Chronic liver disease and cirrhosis (K70,K73-K74)	11	DNR	DNR	4.7 - 18.0
12 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	DNR	DNR	4.5 - 20.3

Table 4 (continued). Leading Causes of Death by Race, Maine, 2009-2018 (page 2 of 2)

Cause of Death	Total Number of Deaths	Crude Rate	Age-adjusted Rate	
			Rate	95% CI
White				
1 Cancer (C00-C97)	32201	251.5	176.1	174.1 - 178.1
2 Heart disease (I00-I09,I11,I13,I20-I51)	27625	215.8	150.4	148.6 - 152.2
3 Chronic lower respiratory disease (J40-J47)	8871	69.3	48.3	47.3 - 49.3
4 Unintentional injury (V01-X59,Y85-Y86)	7196	56.2	49.8	48.6 - 51.0
5 Cerebrovascular disease (I60-I69)	6380	49.8	34.7	33.9 - 35.6
6 Alzheimer's disease (G30)	4994	39.0	26.9	26.1 - 27.6
7 Diabetes (E10-E14)	3916	30.6	21.5	20.8 - 22.2
8 Influenza and pneumonia (J10-J18)	2665	20.8	14.5	13.9 - 15.1
9 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2474	19.3	13.5	13 - 14.1
10 Suicide (X60-X84,Y87.0)	2242	17.5	16.2	15.5 - 16.9
11 Chronic liver disease and cirrhosis (K70,K73-K74)	1664	13.0	9.5	9.0 - 8.0
12 Parkinson's disease (G20-G21)	1562	12.2	8.7	8.3 - 9.1
13 Septicemia (A40-A41)	1330	10.4	7.4	7.0 - 7.8
14 Essential hypertension and hypertensive renal disease (I10,I12,I15)	982	7.7	5.3	4.9 - 5.6
15 Pneumonitis due to solids and liquids (J69)	972	7.6	5.2	4.9 - 5.6

Data Source: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov>. February 2020.

ICD-10 codes are grouped by ICD-10 113 Cause List.

Crude rates are deaths per 100,000 population.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population

95% CI: 95% confidence interval of the rate.

DNR: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

Cause of death categories with numerators less than 10 are suppressed by the WONDER system to protect privacy and are not reported on this table. .

Table 5. Leading Causes of Death by Hispanic Ethnicity, Maine, 2009-2018

Cause of Death	Total Number of Deaths	Crude Rate	Age-adjusted Rate	
			Rate	95% CI
Hispanic				
1 Cancer (C00-C97)	77	39.1	66.8	52 - 84.3
2 Heart disease (I00-I09,I11,I13,I20-I51)	49	24.9	48.8	35.7 - 65.1
3 Unintentional injury (V01-X59,Y85-Y86)	40	20.3	25.0	17.5 - 34.5
4 Diabetes (E10-E14)	20	10.2	17.1	10.3 - 26.7
5 Cerebrovascular disease (I60-I69)	20	10.2	20.4	12.3 - 31.8
6 Suicide (X60-X84,Y87.0)	14	DNR	DNR	3.7 - 12
Non-Hispanic				
1 Cancer (C00-C97)	32390	247.0	175.4	173.5 - 177.4
2 Heart disease (I00-I09,I11,I13,I20-I51)	27715	211.4	149.8	148 - 151.6
3 Chronic lower respiratory disease (J40-J47)	8917	68.0	48.2	47.1 - 49.2
4 Unintentional injury (V01-X59,Y85-Y86)	7270	55.4	49.4	48.2 - 50.6
5 Cerebrovascular disease (I60-I69)	6404	48.8	34.6	33.8 - 35.5
6 Alzheimer's disease (G30)	5000	38.1	26.7	26.0 - 27.5
7 Diabetes (E10-E14)	3948	30.1	21.5	20.8 - 22.2
8 Influenza and pneumonia (J10-J18)	2673	20.4	14.4	13.9 - 15.0
9 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2487	19.0	13.5	12.9 - 14.0
10 Suicide (X60-X84,Y87.0)	2270	17.3	16.1	15.4 - 16.8
11 Chronic liver disease and cirrhosis (K70,K73-K74)	1680	12.8	9.5	9.0 - 9.9
12 Parkinson's disease (G20-G21)	1562	11.9	8.6	8.2 - 9.1
13 Septicemia (A40-A41)	1339	10.2	7.4	7.0 - 7.8
14 Essential hypertension and hypertensive renal disease (I10,I12,I15)	991	7.6	5.3	5.0 - 5.6
15 Pneumonitis due to solids and liquids (J69)	972	7.4	5.2	4.9 - 5.5

Data Source: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov>. February 2020

ICD-10 codes are grouped by ICD-10 113 Cause List.

Crude rates are deaths per 100,000 population.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population

95% CI: 95% confidence interval of the rate.

DNR: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

Cause of death categories with numerators less than 10 are suppressed by the WONDER system to protect privacy and are not reported on this table.

Table 6. Leading Causes of Death by County, Maine, 2018 (page 1 of 5)

Cause of Death	Number of Deaths	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Androscoggin				
1 Cancer (C00-C97)	258	239.6	178.5	156.3 - 200.7
2 Heart disease (I00-I09,I11,I13,I20-I51)	231	214.5	163.4	142.1 - 184.7
3 Alzheimer's disease (G30)	101	93.8	70.4	56.7 - 84.2
4 Unintentional injury (V01-X59,Y85-Y86)	80	74.3	66.3	52.1 - 83.3
5 Chronic lower respiratory disease (J40-J47)	75	85.4	64.7	52.1 - 79.6
6 Cerebrovascular disease (I60-I69)	46	42.7	31.3	22.9 - 41.9
7 Diabetes (E10-E14)	36	33.4	25.7	17.8 - 35.9
8 Influenza and pneumonia (J10-J18)	31	28.8	20.9	14.1 - 29.8
9 Chronic liver disease and cirrhosis (K70,K73-K74)	24	22.3	16.9	10.6 - 25.6
10 Suicide (X60-X84,Y87.0)	19	DNR	DNR	16.0 - 20.7
Aroostook				
1 Cancer (C00-C97)	177	263.7	156.2	131.9 - 180.6
2 Heart disease (I00-I09,I11,I13,I20-I51)	169	251.8	138.9	117.4 - 160.4
3 Chronic lower respiratory disease (J40-J47)	76	113.2	62.9	49.2 - 79.2
4 Cerebrovascular disease (I60-I69)	47	70.0	39.5	28.98 - 52.7
5 Unintentional injury (V01-X59,Y85-Y86)	39	58.1	43.6	29.6 - 61.8
6 Alzheimer's disease (G30)	30	44.7	23.2	15.6 - 33.1
7 Diabetes (E10-E14)	28	41.7	22.9	15.1 - 33.4
8 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	40.2	20.8	13.7 - 30.3
9 Influenza and pneumonia (J10-J18)	15	22.4	12.0	6.7 - 19.8
10 Chronic liver disease and cirrhosis (K70,K73-K74)	15	DNR	DNR	7.8 - 25.0
Cumberland				
1 Cancer (C00-C97)	574	195.5	143.2	131.3 - 155.1
2 Heart disease (I00-I09,I11,I13,I20-I51)	560	190.8	135.5	124.1 - 146.8
3 Unintentional injury (V01-X59,Y85-Y86)	178	60.6	52.5	44.5 - 60.5
4 Chronic lower respiratory disease (J40-J47)	148	50.4	35.2	29.5 - 41.0
5 Cerebrovascular disease (I60-I69)	131	44.6	30.8	25.4 - 36.1
6 Alzheimer's disease (G30)	106	36.1	25.3	20.5 - 30.1
7 Diabetes (E10-E14)	83	28.3	19.5	15.5 - 24.3
8 Influenza and pneumonia (J10-J18)	56	19.1	13.3	10.0 - 17.3
9 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	18.7	12.7	9.6 - 16.6
10 Parkinson's disease (G20-G21)	52	17.7	12.4	9.2 - 16.3

Table 6 (continued). Leading Causes of Death by County, Maine, 2018 (page 2 of 5)

Cause of Death	Number of Deaths	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Franklin				
1 Cancer (C00-C97)	96	321.1	200.5	161.5 - 246.2
2 Heart disease (I00-I09,I11,I13,I20-I51)	58	194.0	125.1	94.5 - 162.5
3 Chronic lower respiratory disease (J40-J47)	25	83.6	44.3	27.4 - 68.7
4 Unintentional injury (V01-X59,Y85-Y86)	23	76.9	68.6	41.3 - 107.2
5 Cerebrovascular disease (I60-I69)	20	DNR	DNR	24.4 - 61.6
6 Diabetes (E10-E14)	15	DNR	DNR	17.4 - 53.5
7 Alzheimer's disease (G30)	15	DNR	DNR	18.3 - 53.9
8 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	DNR	DNR	15.1 - 48.6
9 Influenza and pneumonia (J10-J18)	11	DNR	DNR	12.1 - 50.2
Hancock				
1 Cancer (C00-C97)	142	259.1	157.0	130.0 - 184.0
2 Heart disease (I00-I09,I11,I13,I20-I51)	108	197.0	117.2	94.3 - 140.1
3 Unintentional injury (V01-X59,Y85-Y86)	40	73.0	57.9	39.6 - 81.7
4 Chronic lower respiratory disease (J40-J47)	39	71.2	40.9	28.9 - 56.1
5 Cerebrovascular disease (I60-I69)	27	49.3	26.3	17.3 - 38.3
6 Alzheimer's disease (G30)	18	DNR	DNR	11.0 - 29.4
7 Diabetes (E10-E14)	17	DNR	DNR	11.9 - 35.0
8 Parkinson's disease (G20-G21)	14	DNR	DNR	7.9 - 24.2
9 Influenza and pneumonia (J10-J18)	12	DNR	DNR	5.9 - 19.8
10 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	DNR	DNR	4.6 -17.8
Kennebec				
1 Cancer (C00-C97)	273	223.6	150.9	132.5 - 169.2
2 Heart disease (I00-I09,I11,I13,I20-I51)	273	223.6	152.3	133.9 - 170.7
3 Unintentional injury (V01-X59,Y85-Y86)	102	83.5	75.8	61.0 - 93.1
4 Cerebrovascular disease (I60-I69)	88	72.1	49.8	39.7 - 61.6
5 Chronic lower respiratory disease (J40-J47)	89	72.9	48.1	38.4 - 59.4
6 Diabetes (E10-E14)	62	50.8	34.1	26.0 - 43.9
7 Alzheimer's disease (G30)	48	39.3	26.2	19.3 - 34.7
8 Suicide (X60-X84,Y87.0)	24	19.7	20.0	12.5 - 30.3
9 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	19.7	13.0	8.3 - 19.3
10 Influenza and pneumonia (J10-J18)	23	18.8	12.4	7.8 - 18.5

Table 6 (continued). Leading Causes of Death by County, Maine, 2018 (page 3 of 5)

Cause of Death	Number of Deaths	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Knox				
1 Cancer (C00-C97)	134	336.9	179.9	148.9 - 210.9
2 Heart disease (I00-I09,I11,I13,I20-I51)	112	281.6	154.5	125.5 - 183.4
3 Chronic lower respiratory disease (J40-J47)	32	80.5	44.5	29.8 - 63.9
4 Unintentional injury (V01-X59,Y85-Y86)	27	67.9	50.6	32.1 - 76.0
5 Alzheimer's disease (G30)	21	52.8	28.7	17.8 - 43.9
6 Cerebrovascular disease (I60-I69)	21	52.8	28.4	17.3 - 43.9
7 Diabetes (E10-E14)	10	DNR	DNR	6.9 - 28.7
Lincoln				
1 Heart disease (I00-I09,I11,I13,I20-I51)	98	285.4	153.0	122.5 - 188.7
2 Cancer (C00-C97)	96	279.5	138.4	111.3 - 170.1
3 Cerebrovascular disease (I60-I69)	31	90.3	44.9	30.5 - 63.7
4 Unintentional injury (V01-X59,Y85-Y86)	22	64.1	54.3	31.0 - 88.2
5 Chronic lower respiratory disease (J40-J47)	20	58.2	29.4	18.0 - 45.5
6 Alzheimer's disease (G30)	20	DNR	DNR	18.0 - 45.5
7 Diabetes (E10-E14)	11	DNR	DNR	7.9 - 28.4
Oxford				
1 Cancer (C00-C97)	162	281.2	180.4	151.4 - 209.4
2 Heart disease (I00-I09,I11,I13,I20-I51)	118	204.8	135.5	110.1 - 160.8
3 Chronic lower respiratory disease (J40-J47)	57	98.9	62.4	47.0 - 81.2
4 Unintentional injury (V01-X59,Y85-Y86)	50	86.8	80.4	58.2 - 108.3
5 Diabetes (E10-E14)	30	52.1	32.6	21.9 - 46.9
6 Cerebrovascular disease (I60-I69)	23	39.9	27.2	17.1 - 41.3
7 Alzheimer's disease (G30)	21	36.4	24.3	15.1 - 37.2
8 Influenza and pneumonia (J10-J18)	18	DNR	DNR	11.3 - 31.0
9 Suicide (X60-X84,Y87.0)	12	DNR	DNR	10.0 - 35.7
10 Chronic liver disease and cirrhosis (K70,K73-K74)	11	DNR	DNR	6.0 - 24.8
Penobscot				
1 Cancer (C00-C97)	379	250.8	178.5	160.2 - 196.8
2 Heart disease (I00-I09,I11,I13,I20-I51)	347	229.7	161.3	144.2 - 178.4
3 Unintentional injury (V01-X59,Y85-Y86)	120	79.4	71.1	57.7 - 84.6
4 Chronic lower respiratory disease (J40-J47)	107	70.8	50.4	40.8 - 60.1
5 Cerebrovascular disease (I60-I69)	72	47.7	33.6	26.2 - 42.4
6 Diabetes (E10-E14)	61	40.4	28.4	21.6 - 36.7
7 Alzheimer's disease (G30)	50	33.1	23.5	17.4 - 31.0
8 Suicide (X60-X84,Y87.0)	40	26.5	23.6	16.4 - 32.8

Table 6 (continued). Leading Causes of Death by County, Maine, 2018 (page 4 of 5)

Cause of Death	Number of Deaths	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Penobscot (continued)				
9 Influenza and pneumonia (J10-J18)	38	25.1	17.2	12.1 - 23.8
10 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	18.5	13.6	9.0 - 19.8
Piscataquis				
1 Cancer (C00-C97)	63	375.0	195.6	148.9 - 252.3
2 Heart disease (I00-I09,I11,I13,I20-I51)	58	345.2	179.8	135.5 - 234.0
3 Chronic lower respiratory disease (J40-J47)	20	DNR	DNR	35.6 - 89.9
4 Unintentional injury (V01-X59,Y85-Y86)	18	DNR	DNR	62.9 - 185.4
Sagadahoc				
1 Cancer (C00-C97)	96	269.4	169.0	136.0 - 207.4
2 Heart disease (I00-I09,I11,I13,I20-I51)	86	241.3	157.0	125.6 - 193.9
3 Cerebrovascular disease (I60-I69)	24	67.4	45.6	29.2 - 67.8
4 Unintentional injury (V01-X59,Y85-Y86)	19	DNR	DNR	22.7 - 60.5
5 Chronic lower respiratory disease (J40-J47)	19	DNR	DNR	18.8 - 50.1
6 Suicide (X60-X84,Y87.0)	11	DNR	DNR	16.0 - 61.4
7 Alzheimer's disease (G30)	11	DNR	DNR	9.9 - 35.7
Somerset				
1 Cancer (C00-C97)	138	272.8	180.9	149.6 - 212.2
2 Heart disease (I00-I09,I11,I13,I20-I51)	138	272.8	188.1	156.0 - 220.1
3 Chronic lower respiratory disease (J40-J47)	47	92.9	63.6	46.5 - 84.8
4 Unintentional injury (V01-X59,Y85-Y86)	41	81.0	71.8	50.6 - 99.0
5 Cerebrovascular disease (I60-I69)	18	DNR	DNR	15.2 - 40.5
6 Diabetes (E10-E14)	15	DNR	DNR	10.5 - 32.4
7 Alzheimer's disease (G30)	13	DNR	DNR	9.8 - 31.5
8 Suicide (X60-X84,Y87.0)	13	DNR	DNR	11.7 - 39.7
9 Influenza and pneumonia (J10-J18)	11	DNR	DNR	8.3 - 31.8
Waldo				
1 Cancer (C00-C97)	104	262.0	167.7	135.7 - 205.0
2 Heart disease (I00-I09,I11,I13,I20-I51)	88	221.7	158.3	126.6 - 195.5
3 Chronic lower respiratory disease (J40-J47)	20	50.4	32.3	19.7 - 49.8
4 Cerebrovascular disease (I60-I69)	28	70.5	45.9	30.3 - 66.8
5 Unintentional injury (V01-X59,Y85-Y86)	22	55.4	47.3	28.5 - 73.9
6 Diabetes (E10-E14)	15	DNR	DNR	14.2 - 43.6
7 Alzheimer's disease (G30)	14	DNR	DNR	13.3 - 40.9

Table 6 (continued). Leading Causes of Death by County, Maine, 2018 (page 5 of 5)

Cause of Death	Number of Deaths	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Washington				
1 Heart disease (I00-I09,I11,I13,I20-I51)	105	333.4	198.1	159.3 - 236.9
2 Cancer (C00-C97)	86	273.1	154.4	123.0 - 191.4
3 Unintentional injury (V01-X59,Y85-Y86)	33	104.8	93.1	61.4 - 135.5
4 Chronic lower respiratory disease (J40-J47)	19	60.3	33.1	19.9 - 51.7
5 Cerebrovascular disease (I60-I69)	22	69.9	43.7	27.1 - 66.9
6 Alzheimer's disease (G30)	12	DNR	DNR	11.7 - 39.6
7 Suicide (*U03,X60-X84,Y87.0)	12	DNR	DNR	17.4 - 66.8
8 Diabetes (E10-E14)	12	DNR	DNR	10.3 - 34.7
9 Influenza and pneumonia (J10-J18)	11	DNR	DNR	9.6 - 40.0
10 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	DNR	DNR	9.6 - 34.5
York				
1 Cancer (C00-C97)	494	239.5	157.9	143.6 - 172.1
2 Heart disease (I00-I09,I11,I13,I20-I51)	412	198.8	131.7	118.8 - 144.5
3 Unintentional injury (V01-X59,Y85-Y86)	181	87.8	77.7	65.7 - 89.7
4 Chronic lower respiratory disease (J40-J47)	122	59.2	37.7	30.9 - 44.5
5 Alzheimer's disease (G30)	92	44.6	28.4	22.9 - 34.8
6 Cerebrovascular disease (I60-I69)	84	40.7	26.4	21.0 - 32.8
7 Diabetes (E10-E14)	67	32.5	20.7	16.0 - 26.5
8 Influenza and pneumonia (J10-J18)	50	24.2	16.2	11.9 - 21.4
9 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	21.8	15.0	10.8 - 20.3
10 Suicide (X60-X84,Y87.0)	37	17.9	15.5	10.6 - 21.9

Data Source: Maine CDC, Data Research, and Vital Statistics.

ICD-10 codes are grouped by ICD-10 113 Cause List.

Crude rates are deaths per 100,000 population.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population

95% CI: 95% confidence interval of the rate.

DNR: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

Cause of death categories with numerators less than 10 are suppressed by the WONDER system to protect privacy and are not reported on this table.

Counties with fewer than 10 leading causes are due to small numbers. * ICD codes for terrorism are preceded with '*'.

Table 7. Years of Potential Life Lost (YPLL) before Age 75, by Cause of Death, Maine vs. U.S., 2018

Maine			United States				
Rank	Cause	YPLL	Percent of Total YPLL	Rank	Cause	YPLL	Percent of Total YPLL
	All Causes	91,543			All Causes	21,974,395	
1	Cancer	21,551	23.5%	1	Cancer	4,209,943	19.2%
2	Unintentional Injury	18,631	20.4%	2	Unintentional Injury	3,901,367	17.8%
3	Heart Disease	11,971	13.1%	3	Heart Disease	3,254,225	14.8%
4	Suicide	6,959	7.6%	4	Suicide	1,387,984	6.3%
5	Chronic Lower Respiratory Disease	3,544	3.9%	5	Perinatal Period	801,352	3.6%
7	Diabetes	2,824	3.1%	6	Homicide	758,679	3.5%
6	Liver Disease	2,365	2.6%	7	Diabetes	632,923	2.9%
8	Perinatal Period	2,325	2.5%	8	Liver Disease	632,554	2.9%
9	Cerebrovascular Disease	2,040	2.2%	9	Chronic Lower Respiratory Disease	620,073	2.8%
10	Congenital Anomalies	1,627	1.8%	10	Cerebrovascular Disease	544,130	2.5%
	All Others	17,706	19.3%		All Others	5,231,165	23.8%

Data Source: National Center for Health Statistics (NCHS), National Vital Statistics System. Centers for Disease Control and Prevention: WISQARS Leading Causes of Death. Available at: www.cdc.gov/injury/wisqars

Because numbers are not controlled for population size, YPLL can not be compared across different population groups; only the order of causes can be compared.

Table 8. All Deaths by Year, Maine and U.S, 1999-2018

Year	Maine			U.S.	
	Number of Deaths	Age-adjusted Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
1999	12,261	865.9	850.6 - 881.3	875.6	874.5 - 876.7
2000	12,354	859.4	844.2 - 874.6	869.0	867.9 - 870.1
2001	12,421	848.7	833.7 - 863.6	858.8	857.7 - 859.9
2002	12,694	853.7	838.8 - 868.6	855.9	854.8 - 857.0
2003	12,540	829.8	815.3 - 844.4	843.5	842.4 - 844.5
2004	12,443	813.0	798.6 - 827.3	813.7	812.7 - 814.7
2005	12,868	828.2	813.8 - 842.6	815.0	814.0 - 816.0
2006	12,294	775.1	761.3 - 788.9	791.8	790.8 - 792.8
2007	12,493	771.6	758.0 - 785.3	775.3	774.3 - 776.2
2008	12,541	759.4	746.0 - 772.8	774.9	774.0 - 775.9
2009	12,594	752.6	739.2 - 765.9	749.6	748.6 - 750.5
2010	12,750	749.6	736.4 - 762.8	747.0	746.0 - 747.9
2011	13,001	752.8	739.6 - 766.0	741.3	740.4 - 742.2
2012	12,870	730.4	717.5 - 743.3	732.8	731.9 - 733.7
2013	13,547	754.2	741.2 - 767.2	731.9	731.0 - 732.8
2014	13,510	739.0	726.2 - 751.8	724.6	723.8 - 725.5
2015	14,479	783.5	770.4 - 796.6	733.1	732.2 - 734.0
2016	14,182	759.0	746.1 - 772.0	728.8	727.9 - 729.7
2017	14,676	771.6	758.7 - 784.5	731.9	731.0 - 732.8
2018	14,716	753.9	741.6 - 766.1	723.6	722.8 - 724.5

Data Source for years 1999-2017: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov>. 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.US Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death. 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Feb 13, 2020

All Deaths: Deaths in which any underlying cause of death was coded.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 9. All Deaths by Sex and Year, Maine, 1999-2018

Year	Maine Males			Maine Females		
	Number of Deaths	Age-adjusted Rate		Number of Deaths	Age-adjusted Rate	
		Rate	95% CI		Rate	95% CI
1999	5,788	1,041.4	1,014.0 - 1,068.8	6,473	738.3	720.0 - 756.6
2000	5,877	1,039.7	1,012.6 - 1,066.8	6,477	729.8	711.7 - 747.8
2001	5,928	1,020.9	994.4 - 1,047.3	6,493	720.0	702.2 - 737.8
2002	6,182	1,037.9	1,011.6 - 1,064.2	6,512	715.5	697.9 - 733.2
2003	5,988	986.5	961.0 - 1,011.9	6,552	712.3	694.8 - 729.8
2004	5,885	950.2	925.5 - 974.9	6,558	705.0	687.6 - 722.3
2005	6,286	992.6	967.7 - 1,017.6	6,582	703.0	685.7 - 720.3
2006	6,086	933.5	909.6 - 957.3	6,208	651.8	635.2 - 668.3
2007	6,191	936.0	912.3 - 959.8	6,302	646.5	630.1 - 662.8
2008	6,172	905.7	882.7 - 928.7	6,369	644.5	628.3 - 660.7
2009	6,176	892.5	869.8 - 915.2	6,418	642.8	626.6 - 658.9
2010	6,277	883.6	861.3 - 905.8	6,473	643.8	627.7 - 659.9
2011	6,437	887.8	865.7 - 910.0	6,564	641.9	625.8 - 657.9
2012	6,415	860.8	839.2 - 882.3	6,455	619.9	604.3 - 635.5
2013	6,828	899.8	877.9 - 921.6	6,719	633.9	618.2 - 649.7
2014	6,821	874.1	852.8 - 895.4	6,689	625.3	609.7 - 640.9
2015	7,211	916.4	894.7 - 938.1	7,268	670.7	654.6 - 686.7
2016	7,206	895.3	874.0 - 916.6	6,976	641.6	625.8 - 657.3
2017	7,392	906.1	884.7 - 927.4	7,284	655.6	639.9 - 671.3
2018	7,533	897.3	877.0 - 917.5	7,183	630.5	615.7 - 645.4

Data Source for years 1999-2017: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov>. 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

All Death: Deaths in which any underlying cause of death was coded

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population. 95% CI: 95% confidence interval of the rate.

Table 10. All Deaths by Sex and Age, Maine, 2018

	Number of Deaths	Crude Rate	Age-Adjusted Rate	
		Rate	Rate	95% CI
Maine total	14,716	1,099.5	753.9	741.6 - 766.1
Sex				
Female	7,183	1,051.6	630.5	615.7 - 645.4
Male	7,533	1,149.4	897.3	877.0 - 917.5
Age				
Under 5 yrs	69	107.3	NApp	NApp
05-14 yrs	19	13.5	NApp	NApp
15-24 yrs	96	63.0	NApp	NApp
25-34 yrs	223	139.1	NApp	NApp
35-44 yrs	324	212.6	NApp	NApp
45-54 yrs	720	399.6	NApp	NApp
55-64 yrs	1,745	823.2	NApp	NApp
65-74 yrs	2,777	1,699.9	NApp	NApp
75-84 yrs	3,759	4,745.4	NApp	NApp
85 yrs and over	4,972	14,877.8	NApp	NApp

Data Source: Maine DHHS, CDC, Data Research and Vital Statistics.

All Deaths: Deaths in which any underlying cause of death was coded.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

NApp = Not applicable; age-adjusted rates are not calculated for age-specific groups.

DNR: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

Table 11. All Deaths by Race and Hispanic Ethnicity, Maine, 2009 - 2018

	Average Annual Number	Crude Rate		Age-Adjusted Rate	
		Rate	95% CI	Rate	95% CI
Maine total	13,632	1024.3	1018.9 - 1029.7	754.1	750.5 - 758.1
Race					
American Indian or Alaska Native	66	603.3	557.1 - 649.4	854.9	780.1 - 929.7
Asian or Pacific Islander	36	208.7	187.0 - 230.4	332.6	295.4 - 369.9
Black or African American	55	243.2	222.9 - 263.6	488.2	441.5 - 535.0
White	13,476	1052.5	1046.9 - 1058.2	756.7	752.6 - 760.9
Hispanic Ethnicity					
Non-Hispanic	13,541	1032.7	1027.2 - 1038.2	753.6	749.5 - 757.8
Hispanic or Latino	35	176.2	157.7 - 194.8	293.1	260.5 - 325.7

Data Source: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov>. February 2020.

All Deaths: Deaths in which any underlying cause of death was coded.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 12. All Deaths by County, Maine, 2014 - 2018				
	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	14,302	1072.9	760.8	755.2 - 766.4
County				
Androscoggin	1,140	1060.6	837.4	815.9 - 858.8
Aroostook	875	1283.0	793.8	769.6 - 818.0
Cumberland	2,666	915.2	687.3	675.7 - 698.9
Franklin	346	1153.0	806.2	768.1 - 844.3
Hancock	612	1122.9	714.8	688.9 - 740.8
Kennebec	1,337	1103.5	808.0	788.6 - 827.3
Knox	492	1237.2	731.3	701.5 - 761.1
Lincoln	426	1244.8	711.5	679.5 - 743.5
Oxford	653	1138.3	777.3	750.5 - 804.1
Penobscot	1,632	1072.4	808.5	791.0 - 826.0
Piscataquis	256	1519.5	919.4	866.7 - 972.1
Sagadahoc	371	1052.0	717.6	684.9 - 750.4
Somerset	609	1197.2	868.7	837.7 - 899.6
Waldo	406	1030.1	744.6	712.0 - 777.2
Washington	433	1370.7	896.7	857.8 - 935.6
York	2,046	1008.4	721.2	707.1 - 735.2

Data Source: Maine CDC, Data, Research and Vital Statistics. December 2019

All Deaths: Deaths in which the underlying cause of death was coded as ICD-10 any listed.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 13. Cancer Deaths by Year, Maine and U.S, 1999-2018

Year	Maine			U.S.	
	Number of Deaths	Age-adjusted Rate Rate	Age-adjusted Rate 95% CI	Age-adjusted Rate Rate	Age-adjusted Rate 95% CI
1999	3,035	214.6	207.0 - 222.2	200.8	200.2 - 201.3
2000	3,070	213.6	206.1 - 221.2	199.6	199.1 - 200.2
2001	3,045	208.3	200.9 - 215.7	196.5	195.9 - 197.0
2002	3,206	214.9	207.4 - 222.3	194.3	193.8 - 194.8
2003	3,120	205.4	198.2 - 212.7	190.9	190.4 - 191.4
2004	3,124	202.2	195.1 - 209.3	186.8	186.3 - 187.3
2005	3,218	204.8	197.7 - 211.9	185.1	184.6 - 185.6
2006	3,089	192.7	185.9 - 199.5	181.8	181.3 - 182.3
2007	3,112	189.9	183.2 - 196.6	179.3	178.8 - 179.7
2008	3,093	185.0	178.4 - 191.6	176.4	175.9 - 176.8
2009	3,133	185.2	178.7 - 191.8	173.5	173.1 - 174.0
2010	3,248	187.9	181.3 - 194.4	172.8	172.3 - 173.2
2011	3,201	182.2	175.8 - 188.6	169.0	168.5 - 169.4
2012	3,226	179.3	173.1 - 185.6	166.5	166.1 - 166.9
2013	3,227	175.2	169.0 - 181.3	163.2	162.8 - 163.7
2014	3,209	170.3	164.3 - 176.3	161.2	160.8 - 161.6
2015	3,398	178.0	171.9 - 184.1	158.5	158.1 - 158.9
2016	3,275	168.9	163.0 - 174.9	155.8	155.4 - 156.2
2017	3,391	170.8	164.9 - 176.7	152.5	152.1 - 152.9
2018	3,273	162.0	156.3 - 167.6	149.1	148.7 - 149.5

Maine 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. Data Source for years 1999-2017 (ME + US) and US 2018 Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death. 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Feb 13, 2020

Cancer Deaths: Deaths with cancer as the underlying cause of death.

Cancer: ICD-10 codes C00-C97

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 14. Cancer Deaths by Sex and Year, Maine, 1999-2018

Year	Maine Males			Maine Females		
	Number of Deaths	Age-adjusted Rate Rate	95% CI	Number of Deaths	Age-adjusted Rate Rate	95% CI
1999	1,533	263.0	249.6 - 276.4	1,502	185.9	176.4 - 195.4
2000	1,547	258.7	245.6 - 271.8	1,523	184.1	174.8 - 193.5
2001	1,609	264.7	251.6 - 277.9	1,436	171.3	162.4 - 180.3
2002	1,685	268.7	255.6 - 281.7	1,521	178.9	169.8 - 187.9
2003	1,572	245.5	233.1 - 257.8	1,548	179.8	170.8 - 188.8
2004	1,618	247.6	235.3 - 259.9	1,506	171.2	162.5 - 179.9
2005	1,692	253.4	241.1 - 265.7	1,526	171.5	162.8 - 180.2
2006	1,628	238.1	226.4 - 249.9	1,461	162.1	153.7 - 170.5
2007	1,616	233.1	221.5 - 244.7	1,496	162.6	154.3 - 171.0
2008	1,645	228.8	217.6 - 240.1	1,448	155.4	147.2 - 163.5
2009	1,661	228.3	217.1 - 239.4	1,472	155.6	147.5 - 163.7
2010	1,718	227.1	216.1 - 238.0	1,530	159.8	151.6 - 167.9
2011	1,704	221.2	210.5 - 232.0	1,497	153.6	145.7 - 161.5
2012	1,702	214.1	203.7 - 224.4	1,524	153.4	145.5 - 161.3
2013	1,732	213.9	203.6 - 224.2	1,495	146.6	139.0 - 154.2
2014	1,725	205.5	195.6 - 215.5	1,484	143.1	135.6 - 150.6
2015	1,743	206.4	196.5 - 216.4	1,655	157.3	149.5 - 165.1
2016	1,738	200.4	190.7 - 210.1	1,537	145.7	138.2 - 153.3
2017	1,823	205.6	195.9 - 215.3	1,568	144.9	137.4 - 152.3
2018	1,717	188.8	179.7 - 198.0	1,556	141.1	133.9 - 148.4

Data Source: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov> for 1999-2017. 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Cancer Deaths: Deaths with cancer as the underlying cause of death.

Cancer: ICD-10 codes C00-C97

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 15. Cancer Deaths by Sex and Age, Maine, 2018

	Number of Deaths	Crude Rate		Age-Adjusted Rate	
		Rate	95% CI	Rate	95% CI
Maine total	3,273	244.5	236.1-252.8	162.0	156.3 - 167.6
Sex					
Female	1,556	227.8		141.1	133.9 - 148.4
Male	1,717	262.0		188.8	179.7 - 198.0
Age					
Under 5 yrs	*	*	*	NApp	NApp
05-14 yrs	*	*	*	NApp	NApp
15-24 yrs	*	*	*	NApp	NApp
25-34 yrs	13	DNR	3.7-12.5	NApp	NApp
35-44 yrs	43	28.2	19.8-36.7	NApp	NApp
45-54 yrs	203	112.7	97.2-128.2	NApp	NApp
55-64 yrs	612	288.7	265.8-311.6	NApp	NApp
65-74 yrs	915	560.1	523.8-596.4	NApp	NApp
75-84 yrs	911	1,150.1	1075.4-1224.8	NApp	NApp
85 yrs and over	563	1,684.7	1545.5-1823.8	NApp	NApp

Data Source: Maine DHHS, CDC, Data Research and Vital Statistics.

Cancer Deaths: Deaths with cancer as the underlying cause of death.

Cancer: ICD-10 codes C00-C97

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

NApp = Not applicable; age-adjusted rates are not calculated for age-specific groups.

DNR: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

* = To protect privacy, counts less than 10 are suppressed by CDC Wonder.

Table 16. Cancer Deaths by Race and Hispanic Ethnicity, Maine, 2009 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	3,258	244.8	175.4	173.5 - 177.4
Race				
American Indian or Alaska Native	16	147.1	190.0	156.9 – 223.2
Asian or Pacific Islander	10	58.8	84.6	66.9 – 102.2
Black or African American	12	53.9.7	118.0	95.0 - 141.0
White	3,220	251.5	176.1	174.1 – 178.1
Hispanic Ethnicity				
Hispanic or Latino	8	39.1	66.8	52.0 – 84.3
Non-Hispanic	3,239	247	175.4	173.5 – 177.4

Data Source: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov>
 Cancer Deaths: Deaths with cancer as the underlying cause of death.
 Cancer: ICD-10 codes C00-C97
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 17. Cancer Deaths by County, Maine, 2014 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	3,304	247.9	169.6	167.0 - 172.2
County				
Androscoggin	250	233.0	178.4	168.4 - 188.4
Aroostook	186	272.8	166.8	155.6 - 178.0
Cumberland	599	205.8	153.5	147.9 - 159.1
Franklin	81	270.9	181.0	163.0 - 199.1
Hancock	142	259.6	157.8	145.8 - 169.8
Kennebec	307	253.6	176.5	167.5 - 185.5
Knox	120	300.9	175.0	160.4 - 189.5
Lincoln	100	292.5	156.4	142.0 - 170.8
Oxford	157	274.2	179.4	166.5 - 192.3
Penobscot	356	234.2	171.0	162.9 - 179.1
Piscataquis	66	393.5	215.6	191.8 - 239.5
Sagadahoc	93	264.6	170.2	154.4 - 185.9
Somerset	147	289.4	196.7	182.1 - 211.3
Waldo	97	245.6	165.5	150.3 - 180.8
Washington	103	325.4	197.6	179.8 - 215.3
York	498	245.3	169.5	162.7 - 176.3

Data Source: Maine CDC, Data Research and Vital statistics. March 2020

Cancer Deaths: Deaths with cancer as the underlying cause of death.

Cancer: ICD-10 codes C00-C97

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 18. Heart Disease Deaths by Year, Maine and U.S, 1999-2018

Year	Maine			U.S.	
	Number of Deaths	Age-adjusted Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
1999	3,418	238.7	230.7 - 246.7	266.5	265.8 - 267.1
2000	3,400	233.9	226.0 - 241.7	257.6	257.0 - 258.2
2001	3,272	220.7	213.1 - 228.2	249.5	248.9 - 250.1
2002	3,170	210.6	203.3 - 218.0	244.6	244.1 - 245.2
2003	3,106	203.0	195.9 - 210.2	236.3	235.8 - 236.9
2004	2,948	189.9	183.0 - 196.8	221.6	221.1 - 222.2
2005	2,941	186.5	179.8 - 193.3	216.8	216.3 - 217.4
2006	2,815	174.3	167.8 - 180.8	205.5	205.0 - 206.0
2007	2,852	173.0	166.6 - 179.3	196.1	195.6 - 196.6
2008	2,785	164.6	158.5 - 170.8	192.1	191.6 - 192.6
2009	2,674	156.4	150.4 - 162.3	182.8	182.4 - 183.3
2010	2,628	151.1	145.3 - 156.9	179.1	178.7 - 179.6
2011	2,648	149.5	143.7 - 155.2	173.7	173.3 - 174.2
2012	2,623	145.9	140.2 - 151.6	170.5	170.1 - 170.9
2013	2,807	152.3	146.5 - 158.0	169.8	169.3 - 170.2
2014	2,776	147.9	142.3 - 153.5	167.0	166.6 - 167.4
2015	3,009	157.3	151.6 - 163.0	168.5	168.1 - 168.9
2016	2,907	149.5	144.0 - 155.1	165.5	165.1 - 165.9
2017	2,844	143.5	138.1 - 148.9	165.0	164.6 - 165.4
2018	2,963	146.9	141.6 - 152.3	163.6	163.2 - 164.0

Maine 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. Data Source for years 1999-2017 (ME + US) and US 2018 Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death. 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Feb 13, 2020

Heart Disease Deaths: Deaths with heart disease as the underlying cause of death.

Heart Disease: ICD-10 codes I00-I09, I11, I13, I20-I51.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 19. Heart Disease Deaths by Sex and Year, Maine, 1999-2018

Year	Maine Males			Maine Females		
	Number of Deaths	Age-adjusted Rate	95% CI	Number of Deaths	Age-adjusted Rate	95% CI
1999	1,618	296.7	281.9 - 311.4	1,800	195.9	186.8 - 205.1
2000	1,649	295.4	280.9 - 310.0	1,751	188.1	179.2 - 197.0
2001	1,542	268.6	254.9 - 282.2	1,730	182.3	173.6 - 191.0
2002	1,574	268.2	254.7 - 281.7	1,596	167.3	159.0 - 175.6
2003	1,534	255.2	242.2 - 268.2	1,572	162.0	153.9 - 170.1
2004	1,417	231.9	219.6 - 244.1	1,531	156.4	148.5 - 164.4
2005	1,470	233.6	221.4 - 245.7	1,471	150.1	142.3 - 157.9
2006	1,470	226.2	214.5 - 238.0	1,345	133.4	126.2 - 140.7
2007	1,504	228.2	216.5 - 240.0	1,348	131.5	124.3 - 138.6
2008	1,413	205.5	194.6 - 216.4	1,372	129.9	122.9 - 136.9
2009	1,376	200.5	189.8 - 211.3	1,298	123.2	116.4 - 130.0
2010	1,362	193.6	183.1 - 204.0	1,266	119.0	112.3 - 125.7
2011	1,374	189.6	179.4 - 199.8	1,274	117.1	110.5 - 123.7
2012	1,377	185.5	175.5 - 195.5	1,246	113.6	107.1 - 120.1
2013	1,476	193.7	183.6 - 203.8	1,331	119.5	112.9 - 126.1
2014	1,439	184.2	174.5 - 193.9	1,337	118.1	111.7 - 124.6
2015	1,602	200.3	190.3 - 210.3	1,407	121.9	115.3 - 128.5
2016	1,592	193.6	183.8 - 203.4	1,315	112.8	106.5 - 119.1
2017	1,533	183.3	173.9 - 192.7	1,311	111.6	105.3 - 117.8
2018	1,648	193.7	184.2 - 203.22	1,315	110.0	103.9 - 116.11

Data Source: 1999-2017: Detailed Mortality Files accessed through CDC Wonder.

<https://wonder.cdc.gov> 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Heart Disease Deaths: Deaths with heart disease as the underlying cause of death.

Heart Disease: ICD-10 codes I00-I09, I11, I13, I20-I51.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 20. Heart Disease Deaths by Sex and Age, Maine, 2018

	Number of Deaths	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	2,963	221.4	146.9	141.6 - 152.3
Sex				
Female	1,315	192.5	110.0	103.9 - 116.1
Male	1,648	251.5	193.7	184.2 - 203.2
Age				
Under 5 yrs	*	*	NApp	NApp
05-14 yrs	*	*	NApp	NApp
15-24 yrs	*	*	NApp	NApp
25-34 yrs	*	*	NApp	NApp
35-44 yrs	43	28.2	NApp	NApp
45-54 yrs	112	62.2	NApp	NApp
55-64 yrs	285	134.4	NApp	NApp
65-74 yrs	497	304.2	NApp	NApp
75-84 yrs	732	924.1	NApp	NApp
85 yrs and over	1,278	3,824.2	NApp	NApp

Data Source: Maine DHHS, CDC, Data Research and Vital Statistics.

Heart Disease Deaths: Deaths with heart disease as the underlying cause of death.

Heart Disease: ICD-10 codes I00-I09, I11, I13, I20-I51.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

NApp = Not applicable; age-adjusted rates are not calculated for age-specific groups.

* = To protect privacy, counts less than 10 are suppressed.

Table 21. Heart Disease Deaths by Race and Hispanic Ethnicity, Maine, 2009 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	2,788	209.5	149.9	148.1 - 181.7
Race				
American Indian or Alaska Native	10	95.6	163.1	127.5 - 198.6
Asian or Pacific Islander	6	34.7	62.1	46.2 - 81.6
Black or African American	9	41.1	67.9	77.4 - 122.2
White	2,763	215.8	150.4	148.6 - 152.2
Hispanic Ethnicity				
Hispanic or Latino	5	24.9	48.8	35.7 - 65.1
Non-Hispanic	2,772	211.4	149.8	148 - 151.6

Data Source: Detailed Mortality Files accessed through CDC Wonder.
<https://wonder.cdc.gov>
Heart Disease Deaths: Deaths with heart disease as the underlying cause of death.
Heart Disease: ICD-10 codes I00-I09, I11, I13, I20-I51.
Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
95% CI: 95% confidence interval of the rate.
Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 22. Heart Disease Deaths by County, Maine, 2014 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	2,889	216.8	148.5	146.1 - 151.0
County				
Androscoggin	231	215.0	166.0	156.4 - 175.6
Aroostook	187	274.8	160.0	149.5 - 170.5
Cumberland	501	172.0	124.8	119.9 - 129.8
Franklin	68	225.7	152.8	136.3 - 169.2
Hancock	126	231.0	142.8	131.4 - 154.2
Kennebec	270	222.5	158.7	150.2 - 167.3
Knox	112	280.7	153.3	140.4 - 166.2
Lincoln	92	270.3	147.2	133.2 - 161.2
Oxford	119	206.9	138.5	127.2 - 149.8
Penobscot	334	219.7	161.1	153.3 - 168.9
Piscataquis	56	333.1	190.4	167.3 - 213.5
Sagadahoc	80	225.5	147.9	133.3 - 162.5
Somerset	140	274.4	195.1	180.5 - 209.7
Waldo	94	239.5	169.9	154.4 - 185.4
Washington	91	288.7	179.3	162.5 - 196.2
York	388	191.4	131.6	125.7 - 137.5

Data Source: Maine CDC, Data, Research and Vital Statistics. December 2019.

Heart Disease Deaths: Deaths with heart disease as the underlying cause of death.

Heart Disease: ICD-10 codes I00-I09, I11, I13, I20-I51.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 23. Chronic Lower Respiratory Disease Deaths by Year, Maine and U.S, 1999-2018

Year	Maine			U.S.	
	Number of Deaths	Age-adjusted Rate	95% CI	Age-adjusted Rate	95% CI
1999	751	52.6	48.8 - 56.3	45.4	45.2 - 45.7
2000	768	53.0	49.2 - 56.7	44.2	43.9 - 44.4
2001	800	54.3	50.6 - 58.1	43.9	43.6 - 44.1
2002	791	52.9	49.2 - 56.6	43.9	43.6 - 44.1
2003	782	51.3	47.7 - 54.9	43.7	43.5 - 44.0
2004	766	49.7	46.2 - 53.2	41.6	41.4 - 41.8
2005	830	53.2	49.6 - 56.8	43.9	43.7 - 44.1
2006	783	49.1	45.6 - 52.5	41.0	40.8 - 41.2
2007	728	44.5	41.2 - 47.7	41.4	41.1 - 41.6
2008	794	47.8	44.4 - 51.1	44.7	44.4 - 44.9
2009	821	48.6	45.3 - 52.0	42.7	42.4 - 42.9
2010	809	47.4	44.1 - 50.7	42.2	42.0 - 42.4
2011	842	48.4	45.1 - 51.7	42.5	42.3 - 42.7
2012	817	45.4	42.2 - 48.5	41.5	41.3 - 41.8
2013	902	49.1	45.8 - 52.3	42.1	41.9 - 42.4
2014	896	48.1	44.9 - 51.3	40.5	40.2 - 40.7
2015	1,025	53.6	50.3 - 56.9	41.6	41.4 - 41.8
2016	928	47.4	44.3 - 50.5	40.6	40.4 - 40.8
2017	982	48.7	45.6 - 52.8	40.9	40.7 - 41.1
2018	938	45.6	42.6 - 48.5	39.7	39.5 - 39.9

Research and Vital Statistics. Data Source for years 1999-2017 (ME + US) and US 2018 Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death. 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Chronic Lower Respiratory Disease (CLRD) Deaths: Deaths with CLRD as the underlying cause of death.
 Chronic Lower Respiratory Disease: ICD-10 codes J40-J47.
 Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.

Table 24. Chronic Lower Respiratory Disease Deaths by Sex and Year, Maine, 1999-2018

Year	Maine Males			Maine Females		
	Number of Deaths	Age-adjusted Rate	95% CI	Number of Deaths	Age-adjusted Rate	95% CI
1999	363	66.4	59.5 - 73.4	388	44.7	40.2 - 49.2
2000	375	67.5	60.5 - 74.4	393	45.5	41.0 - 50.0
2001	367	64.0	57.3 - 70.6	433	49.6	44.8 - 54.3
2002	418	70.4	63.5 - 77.2	373	41.5	37.3 - 45.8
2003	353	58.8	52.6 - 65.0	429	47.9	43.3 - 52.4
2004	371	59.9	53.7 - 66.1	395	43.5	39.1 - 47.8
2005	402	63.6	57.3 - 70.0	428	46.4	42.0 - 50.9
2006	390	60.3	54.3 - 66.4	393	42.1	37.9 - 46.3
2007	336	50.7	45.2 - 56.1	392	40.6	36.5 - 44.6
2008	384	56.3	50.6 - 62.0	410	42.4	38.3 - 46.6
2009	387	56.4	50.7 - 62.1	434	44.1	39.9 - 48.3
2010	376	53.3	47.9 - 58.8	433	44.0	39.8 - 48.2
2011	400	54.4	49.0 - 59.8	442	44.2	40.0 - 48.4
2012	383	50.4	45.2 - 55.5	434	42.4	38.4 - 46.5
2013	454	58.5	53.0 - 63.9	448	42.4	38.4 - 46.4
2014	459	56.8	51.5 - 62.1	437	42.1	38.1 - 46.1
2015	471	58.0	52.6 - 63.3	554	50.9	46.6 - 55.2
2016	467	55.4	50.3 - 60.5	461	42.0	38.0 - 45.9
2017	475	54.3	49.3 - 59.3	507	44.7	40.7 - 48.7
2018	476	53.5	48.6 - 58.4	462	39.7	36.0 - 43.3

Data Source for years 1999-2017: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov> . 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Chronic Lower Respiratory Disease (CLRD) Deaths: Deaths with CLRD as the underlying cause of death.

Chronic Lower Respiratory Disease: ICD-10 codes J40-J47.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 25. Chronic Lower Respiratory Disease Deaths by Sex and Age, Maine, 2018

	Number of Deaths	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	938	70.1	45.6	42.6-48.5
Sex				
Female	462	67.6	39.7	36.0 - 43.3
Male	476	72.6	53.5	48.6 - 58.4
Age				
Under 5 yrs	*	*	NApp	NApp
05-14 yrs	*	*	NApp	NApp
15-24 yrs	*	*	NApp	NApp
25-34 yrs	*	*	NApp	NApp
35-44 yrs	*	*	NApp	NApp
45-54 yrs	19	10.5	NApp	NApp
55-64 yrs	120	56.6	NApp	NApp
65-74 yrs	220	134.7	NApp	NApp
75-84 yrs	311	392.6	NApp	NApp
85 yrs and over	263	787.0	NApp	NApp

Data Source: Maine DHHS, CDC, Data Research and Vital Statistics.

Chronic Lower Respiratory Disease (CLRD) Deaths: Deaths with CLRD as the underlying cause of death.

Chronic Lower Respiratory Disease: ICD-10 codes J40-J47.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

NApp = Not applicable; age-adjusted rates are not calculated for age-specific groups.

* = To protect privacy, counts less than 10 are suppressed.

Table 26. Chronic Lower Respiratory Disease Deaths by Race and Hispanic Ethnicity, Maine, 2009 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	896	67.0	48.0	47.1 - 49.2
Race				
American Indian or Alaska Native	7	61.6	85.0	64 - 110.6
Asian or Pacific Islander	*	*	*	*
Black or African American	1	^	^	9.3 - 31.5
White	887	69.3	48.3	47.3 - 49.3
Hispanic Ethnicity				
Hispanic or Latino	*	*	*	*
Non-Hispanic	878	66.9	48.5	47.4 - 49.5

Data Source: Detailed Mortality Files accessed through CDC Wonder.

<https://wonder.cdc.gov>. *

Chronic Lower Respiratory Disease (CLRD) Deaths: Deaths with CLRD as the underlying cause of death.

Chronic Lower Respiratory Disease: ICD-10 codes J40-J47.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

DNR: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

***Totals are not available due to suppression constraints

* = To protect privacy, counts less than 10 are suppressed by CDC Wonder.

^ =Death rates are flagged by CDC WONDER as Unreliable when the rate is calculated with a numerator of 20 or less

**Table 27. Chronic Lower Respiratory
Disease Deaths by County, Maine, 2014 - 2018**

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	951	71.4	48.5	47.1 - 49.9
County				
Androscoggin	80	74.1	57.4	51.7 - 63.1
Aroostook	69	101.8	59.2	52.9 - 65.6
Cumberland	154	52.8	38.4	35.6 - 41.1
Franklin	25	84.5	55.1	45.4 - 64.8
Hancock	37	68.2	41.8	35.6 - 47.9
Kennebec	89	73.8	51.7	46.9 - 56.6
Knox	30	76.5	44.5	37.1 - 51.9
Lincoln	24	69.0	35.5	29.0 - 42.0
Oxford	58	101.5	65.9	58.2 - 73.6
Penobscot	115	75.7	56.1	51.5 - 60.8
Piscataquis	17	103.1	56.3	44.9 - 69.8
Sagadahoc	22	62.9	40.0	32.5 - 47.6
Somerset	50	99.1	69.0	60.3 - 77.6
Waldo	25	63.4	42.0	34.5 - 49.5
Washington	27	86.7	53.3	44.1 - 62.5
York	126	62.4	42.1	38.9 - 45.5

Data Source: Maine CDC, Data, Research and Vital Statistics. December 2019

Chronic Lower Respiratory Disease (CLRD) Deaths: Deaths with CLRD as the underlying cause of death.

Chronic Lower Respiratory Disease: ICD-10 codes J40-J47.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 28. Unintentional Injury Deaths by Year, Maine and U.S., 1999-2018

Year	Maine			U.S.	
	Number of Deaths	Age-adjusted Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
1999	458	34.9	31.7 - 38.1	35.3	35.1 - 35.5
2000	407	30.9	27.9 - 33.9	34.9	34.7 - 35.2
2001	490	36.3	33.1 - 39.6	35.7	35.5 - 35.9
2002	511	38.1	34.7 - 41.4	37.1	36.9 - 37.3
2003	516	38.0	34.7 - 41.3	37.6	37.4 - 37.8
2004	480	35.5	32.3 - 38.8	38.1	37.8 - 38.3
2005	579	41.7	38.3 - 45.1	39.5	39.3 - 39.7
2006	572	40.8	37.4 - 44.2	40.2	40.0 - 40.5
2007	584	41.6	38.1 - 45.0	40.4	40.1 - 40.6
2008	628	43.4	39.9 - 46.9	39.2	39.0 - 39.5
2009	602	40.9	37.6 - 44.3	37.5	37.3 - 37.7
2010	540	36.5	33.4 - 39.7	38.0	37.8 - 38.2
2011	574	38.3	35.1 - 41.6	39.1	38.9 - 39.4
2012	602	40.4	37.0 - 43.7	39.1	38.8 - 39.3
2013	644	42.6	39.2 - 46.1	39.4	39.2 - 39.6
2014	690	45.8	42.2 - 49.4	40.5	40.3 - 40.7
2015	802	53.8	49.9 - 57.7	43.2	43.0 - 43.4
2016	909	62.4	58.1 - 66.6	47.4	47.2 - 47.6
2017	990	68.0	63.6 - 72.5	49.4	49.2 - 49.6
2018	996	64.7	60.4 - 68.9	48.0	47.8 - 48.2

Maine 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. Data Source for years 1999-2017 (ME + US) and US 2018 Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death. 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Feb 13, 2020

Unintentional Injury Deaths: Deaths with unintentional injury as the underlying cause of death.

Unintentional Injury: ICD-10 codes V01-X59, Y85-Y86.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 29. Unintentional Injury Deaths by Sex and Year, Maine, 1999-2018

Year	Maine Males			Maine Females		
	Number of Deaths	Age-adjusted Rate Rate	95% CI	Number of Deaths	Age-adjusted Rate Rate	95% CI
1999	309	53.3	47.2 - 59.3	149	20.1	16.8 - 23.4
2000	264	44.0	38.7 - 49.4	143	18.6	15.5 - 21.8
2001	305	50.4	44.7 - 56.1	185	24.0	20.4 - 27.5
2002	313	51.1	45.4 - 56.8	198	26.5	22.7 - 30.4
2003	334	53.7	47.9 - 59.6	182	23.8	20.3 - 27.4
2004	293	47.9	42.4 - 53.5	187	24.3	20.7 - 27.9
2005	372	59.1	53.0 - 65.2	207	25.7	22.1 - 29.3
2006	371	56.9	51.1 - 62.8	201	25.2	21.6 - 28.8
2007	381	59.6	53.5 - 65.7	203	25.4	21.7 - 29.0
2008	386	58.7	52.8 - 64.7	242	29.3	25.5 - 33.2
2009	384	57.9	52.0 - 63.8	218	25.1	21.6 - 28.6
2010	345	51.9	46.3 - 57.5	195	22.8	19.4 - 26.1
2011	354	52.5	46.9 - 58.1	220	25.0	21.5 - 28.6
2012	372	56.7	50.8 - 62.6	230	25.0	21.6 - 28.5
2013	405	59.0	53.1 - 65.0	239	27.5	23.7 - 31.2
2014	429	62.4	56.3 - 68.6	261	30.1	26.2 - 34.1
2015	501	74.0	67.3 - 80.7	301	34.8	30.6 - 39.0
2016	555	83.6	76.4 - 90.8	354	41.0	36.3 - 45.6
2017	631	95.4	87.6 - 103.1	359	41.4	36.7 - 46.0
2018	615	89.9	82.5 - 97.3	381	40.7	36.3 - 45.2

Data Source for years 1999-2017: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov> . 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Unintentional Injury Deaths: Deaths with unintentional injury as the underlying cause of death.

Unintentional Injury: ICD-10 codes V01-X59,Y85-Y86.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

**Table 30. Unintentional Injury
Deaths by Sex and Age, Maine, 2018**

	Number of Deaths	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	996	74.4	64.7	60.4 - 68.9
Sex				
Female	381	55.8	40.7	36.3-45.2
Male	615	93.8	89.9	82.5-97.3
Age				
Under 5 yrs	*	*	NApp	NApp
05-14 yrs	*	*	NApp	NApp
15-24 yrs	39	25.6	NApp	NApp
25-34 yrs	119	74.2	NApp	NApp
35-44 yrs	126	82.7	NApp	NApp
45-54 yrs	115	63.8	NApp	NApp
55-64 yrs	116	54.7	NApp	NApp
65-74 yrs	86	52.6	NApp	NApp
75-84 yrs	139	175.5	NApp	NApp
85 yrs and over	239	715.2	NApp	NApp

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Unintentional Injury Deaths: Deaths with unintentional injury as the underlying cause of death.

Unintentional Injury: ICD-10 codes V01-X59,Y85-Y86.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

NApp = Not applicable; age-adjusted rates are not calculated for age-specific groups.

* = To protect privacy, counts less than 10 are suppressed

Table 31. Unintentional Injury Deaths by Race and Hispanic Ethnicity, Maine, 2009 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	735	55.2	49.4	48.2 - 50.6
Race				
American Indian or Alaska Native	7	65.3	70.9	54.1 - 91.3
Asian or Pacific Islander	3	15.9	21.8	13.7 - 33.1
Black or African American	6	26.0	33.2	24.1 - 44.5
White	720	56.2	49.8	48.6 - 51.0
Hispanic Ethnicity				
Hispanic or Latino	4	20.3	25.0	17.5 - 34.5
Non-Hispanic	727	55.4	49.4	48.2 - 50.6

Data Source: Detailed Mortality Files accessed through CDC Wonder.
<https://wonder.cdc.gov>

Unintentional Injury Deaths: Deaths with unintentional injury as the underlying cause of death.

Unintentional Injury: ICD-10 codes V01-X59,Y85-Y86.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 32. Unintentional Injury Deaths by County, Maine, 2014 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	874	65.6	58.8	57.0 - 60.7
County				
Androscoggin	66	61.0	56.9	50.6 - 63.3
Aroostook	38	55.7	47.3	40.0 - 54.7
Cumberland	173	59.5	52.8	49.2 - 56.5
Franklin	18	61.2	55.4	43.9 - 68.9
Hancock	37	68.2	59.0	49.8 - 68.2
Kennebec	82	67.7	62.2	55.9 - 68.5
Knox	30	76.0	60.3	55.9 - 68.5
Lincoln	23	68.4	58.4	47.0 - 71.7
Oxford	34	58.6	51.9	43.5 - 60.4
Penobscot	99	64.8	60.0	54.5 - 65.6
Piscataquis	15	91.3	82.5	63.3 - 105.8
Sagadahoc	18	50.4	41.1	32.5 - 51.2
Somerset	37	71.9	66.7	56.5 - 76.9
Waldo	25	64.4	59.4	48.4 - 70.3
Washington	32	101.3	98.6	82.0 - 115.1
York	146	71.8	65.6	60.6 - 70.6

Data Source: Maine CDC, Data, Research and Vital Statistics. December 2019. Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Unintentional Injury Deaths: Deaths with unintentional injury as the underlying cause of death.

Unintentional Injury: ICD-10 codes V01-X59, Y85-Y86.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 33. Cerebrovascular Disease Deaths by Year, Maine and U.S, 1999-2018

Year	Maine			U.S.	
	Number of Deaths	Age-adjusted Rate	95% CI	Age-adjusted Rate	95% CI
1999	879	61.2	57.1 - 65.2	61.6	61.3 - 61.9
2000	833	57.0	53.1 - 60.9	60.9	60.6 - 61.2
2001	822	55.2	51.4 - 59.0	58.4	58.1 - 58.7
2002	823	54.5	50.7 - 58.2	57.2	57.0 - 57.5
2003	802	52.2	48.6 - 55.8	54.6	54.3 - 54.8
2004	801	51.5	47.9 - 55.1	51.2	50.9 - 51.4
2005	693	43.8	40.5 - 47.1	48.0	47.7 - 48.2
2006	670	41.5	38.3 - 44.7	44.8	44.6 - 45.0
2007	664	40.4	37.3 - 43.5	43.5	43.3 - 43.7
2008	671	40.4	37.3 - 43.5	42.1	41.8 - 42.3
2009	647	38.0	35.1 - 41.0	39.6	39.4 - 39.8
2010	602	34.5	31.7 - 37.3	39.1	38.9 - 39.3
2011	630	35.8	33.0 - 38.6	37.9	37.7 - 38.1
2012	619	34.3	31.5 - 37.0	36.9	36.7 - 37.1
2013	620	33.4	30.7 - 36.0	36.2	36.0 - 36.4
2014	628	33.2	30.6 - 35.9	36.5	36.3 - 36.7
2015	616	32.6	30.0 - 35.2	37.6	37.4 - 37.8
2016	663	34.4	31.7 - 37.1	37.3	37.1 - 37.5
2017	736	37.5	34.7 - 40.2	37.6	37.4 - 37.8
2018	691	33.8	31.2 - 36.3	37.1	36.9 - 37.3

Maine 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. Data Source for years 1999-2017 (ME + US) and US 2018 Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death. 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Feb 13, 2020

Cerebrovascular Disease Deaths: Deaths with cerebrovascular disease as the underlying cause of death.

Cerebrovascular Disease: ICD-10 codes I60-I69.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 34. Cerebrovascular Disease Deaths by Sex and Year, Maine, 1999-2018

Year	Maine Males			Maine Females		
	Number of Deaths	Age-adjusted Rate		Number of Deaths	Age-adjusted Rate	
		Rate	95% CI		Rate	95% CI
1999	339	64.6	57.6 - 71.6	540	57.5	52.6 - 62.4
2000	310	59.0	52.3 - 65.7	523	55.2	50.4 - 60.0
2001	355	64.2	57.4 - 71.0	467	48.2	43.8 - 52.6
2002	339	61.2	54.5 - 67.8	484	49.7	45.2 - 54.2
2003	307	54.2	48.0 - 60.3	495	50.9	46.3 - 55.4
2004	304	51.6	45.7 - 57.4	497	50.1	45.6 - 54.5
2005	262	43.4	38.1 - 48.7	431	43.3	39.2 - 47.4
2006	252	40.3	35.2 - 45.3	418	41.4	37.4 - 45.5
2007	240	38.1	33.2 - 43.0	424	41.0	37.0 - 44.9
2008	268	41.5	36.5 - 46.5	403	39.3	35.4 - 43.3
2009	244	36.8	32.2 - 41.5	403	38.1	34.3 - 41.9
2010	243	35.2	30.7 - 39.7	359	33.6	30.0 - 37.1
2011	239	32.9	28.6 - 37.1	391	37.1	33.3 - 40.8
2012	227	30.8	26.8 - 34.9	392	35.2	31.7 - 38.8
2013	245	32.6	28.4 - 36.7	375	33.4	29.9 - 36.8
2014	276	35.9	31.6 - 40.2	352	30.9	27.6 - 34.2
2015	230	29.7	25.8 - 33.6	386	34.1	30.6 - 37.6
2016	261	32.9	28.8 - 37.0	402	35.3	31.8 - 38.9
2017	265	32.4	28.4 - 36.3	471	40.8	37.0 - 44.6
2018	275	32.0	28.1 - 35.8	416	34.1	30.7 - 37.4

Data Source for years 1999-2017: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov> . 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Cerebrovascular Disease Deaths: Deaths with cerebrovascular disease as the underlying cause of death.

Cerebrovascular Disease: ICD-10 codes I60-I69.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 35. Cerebrovascular Disease Deaths by Sex and Age, Maine, 2018

	Number of Deaths	Crude Rate		Age-Adjusted Rate	
		Rate	95% CI	Rate	95% CI
Maine total	691	51.6	47.8 - 55.5	33.8	31.2 - 36.3
Sex					
Female	416	60.9		34.1	30.7 - 37.4
Male	275	42.0		32.0	28.1 - 35.8
Age					
Under 5 yrs	*	*	*	NApp	NApp
05-14 yrs	*	*	*	NApp	NApp
15-24 yrs	*	*	*	NApp	NApp
25-34 yrs	*	*	*	NApp	NApp
35-44 yrs	*	*	*	NApp	NApp
45-54 yrs	13	DNR	3.3 - 11.1	NApp	NApp
55-64 yrs	59	27.8	20.7 - 34.9	NApp	NApp
65-74 yrs	101	61.8	49.8 - 73.9	NApp	NApp
75-84 yrs	182	229.8	196.4 - 263.1	NApp	NApp
85 yrs and over	329	984.5	878.1 - 1090.9	NApp	NApp

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.
 Cerebrovascular Disease Deaths: Deaths with cerebrovascular disease as the underlying cause of death.
 Cerebrovascular Disease: ICD-10 codes I60-I69.
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.
 NApp = Not applicable; age-adjusted rates are not calculated for age-specific groups.
 * = To protect privacy, counts less than 10 are suppressed.

Table 36. Cerebrovascular Disease Deaths by Race and Hispanic Ethnicity, Maine, 2009 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	645	48.5	34.7	33.8 - 35.6
Race				
American Indian or Alaska Native	2	18.4	37.6	21.9 - 60.1
Asian or Pacific Islander	3	18.8	34.2	23.1 - 48.8
Black or African American	2	^	^	12.1 - 31.5
White	638	49.8	34.7	33.9 - 35.6
Hispanic Ethnicity				
Hispanic or Latino	2	10.2	20.4	12.3 - 31.8
Non-Hispanic	640	48.8	34.6	33.8 - 35.5

Data Source: Detailed Mortality Files accessed through CDC Wonder.

<https://wonder.cdc.gov>

Cerebrovascular Disease Deaths: Deaths with cerebrovascular disease as the underlying cause of death.

Cerebrovascular Disease: ICD-10 codes I60-I69

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

^: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

* = To protect privacy, counts less than 10 are suppressed by CDC Wonder.

Table 37. Cerebrovascular Disease Deaths by County, Maine, 2014 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	664	49.8	34.2	33.0 - 35.4
County				
Androscoggin	54	49.9	38.7	34.0 - 43.4
Aroostook	41	59.8	34.9	30.0 - 39.7
Cumberland	127	43.7	31.8	29.3 - 34.3
Franklin	18	60.6	39.9	32.1 - 49.1
Hancock	31	57.6	34.3	28.8 - 39.8
Kennebec	69	57.3	40.9	36.5 - 45.2
Knox	22	55.8	30.4	24.6 - 36.3
Lincoln	28	82.5	42.3	35.2 - 49.5
Oxford	24	42.2	28.1	23.0 - 33.1
Penobscot	71	46.9	34.5	30.9 - 38.1
Piscataquis	12	69.9	40.9	30.7 - 53.6
Sagadahoc	24	68.0	45.7	37.4 - 54.0
Somerset	24	46.4	33.8	27.7 - 40.0
Waldo	21	52.3	37.4	30.4 - 45.6
Washington	16	51.3	32.3	25.5 - 40.4
York	81	39.9	27.7	25.0 - 30.4

Data Source: Maine CDC: Data, Research and Vital Statistics. December 2019.
 Cerebrovascular Disease Deaths: Deaths with cerebrovascular disease as the underlying cause of death.
 Cerebrovascular Disease: ICD-10 codes I60-I69
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 38. Alzheimer's Disease Deaths by Year, Maine and U.S., 1999-2018

Year	Maine			U.S.	
	Number of Deaths	Age-adjusted Rate Rate	Age-adjusted Rate 95% CI	Age-adjusted Rate Rate	Age-adjusted Rate 95% CI
1999	429	29.6	26.8 - 32.4	16.5	16.3 - 16.6
2000	466	31.6	28.7 - 34.5	18.1	17.9 - 18.2
2001	471	31.3	28.5 - 34.1	19.3	19.1 - 19.5
2002	513	33.6	30.7 - 36.5	20.8	20.6 - 21.0
2003	467	30.2	27.5 - 32.9	22.1	21.9 - 22.2
2004	512	32.7	29.9 - 35.5	22.6	22.4 - 22.8
2005	476	29.9	27.2 - 32.6	24.0	23.8 - 24.2
2006	477	29.2	26.6 - 31.8	23.7	23.5 - 23.9
2007	470	28.1	25.6 - 30.7	23.8	23.7 - 24.0
2008	450	26.3	23.9 - 28.8	25.8	25.6 - 26.0
2009	466	26.9	24.5 - 29.4	24.2	24.0 - 24.3
2010	502	28.5	26.0 - 31.1	25.1	24.9 - 25.3
2011	481	26.8	24.4 - 29.2	24.7	24.6 - 24.9
2012	432	23.5	21.2 - 25.7	23.8	23.6 - 23.9
2013	401	21.6	19.4 - 23.7	23.5	23.4 - 23.7
2014	434	22.7	20.5 - 24.8	25.4	25.3 - 25.6
2015	544	28.4	26.0 - 30.8	29.4	29.3 - 29.6
2016	577	29.6	27.2 - 32.1	30.3	30.1 - 30.5
2017	601	30.4	27.9 - 32.8	31.0	30.9 - 31.2
2018	580	28.4	26.0 - 30.7	30.5	30.3 - 30.7

Maine 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. Data Source for years 1999-2017 (ME + US) and US 2018 Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death. 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Feb 13, 2020

Alzheimer's Disease Deaths: Deaths with Alzheimer's disease as the underlying cause of death.

Alzheimer's Disease: ICD-10 code G30.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 39. Alzheimer's Disease Deaths by Sex and Year, Maine, 1999-2018

Year	Maine Males			Maine Females		
	Number of Deaths	Age-adjusted Rate	95% CI	Number of Deaths	Age-adjusted Rate	95% CI
1999	107	22.1	17.9 - 26.4	322	33.1	29.5 - 36.7
2000	143	29.2	24.3 - 34.1	323	32.6	29.0 - 36.2
2001	121	24.2	19.8 - 28.6	350	34.7	31.1 - 38.4
2002	129	24.9	20.5 - 29.2	384	37.9	34.1 - 41.7
2003	148	28.2	23.6 - 32.8	319	31.4	27.9 - 34.9
2004	123	22.5	18.5 - 26.6	389	37.9	34.1 - 41.7
2005	135	24.6	20.4 - 28.8	341	33.0	29.5 - 36.5
2006	133	23.0	19.1 - 26.9	344	32.2	28.8 - 35.6
2007	125	21.1	17.4 - 24.8	345	31.9	28.5 - 35.3
2008	130	21.6	17.9 - 25.4	320	29.2	25.9 - 32.4
2009	147	23.7	19.9 - 27.6	319	28.8	25.6 - 32.0
2010	151	23.7	19.9 - 27.5	351	31.3	28.0 - 34.6
2011	156	23.6	19.9 - 27.3	325	28.5	25.3 - 31.6
2012	129	19.1	15.8 - 22.5	303	26.2	23.2 - 29.2
2013	110	15.7	12.8 - 18.7	291	25.2	22.3 - 28.2
2014	135	18.7	15.5 - 21.9	299	25.1	22.2 - 28.0
2015	168	23.2	19.6 - 26.7	376	31.8	28.5 - 35.1
2016	155	20.7	17.4 - 24.0	422	34.9	31.5 - 38.3
2017	156	20.0	16.9 - 23.2	445	36.6	33.1 - 40.0
2018	164	20.4	17.3 - 23.5	416	33.2	29.9 - 36.4

Data Source for years 1999-2017: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov> . 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Alzheimer's Disease Deaths: Deaths with Alzheimer's disease as the underlying cause of death.
Alzheimer's Disease: ICD-10 code G30.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 40. Alzheimer's Disease Deaths by Sex and Age, Maine, 2018					
	Number of Deaths	Crude Rate		Age-Adjusted Rate	
		Rate	95% CI	Rate	95% CI
Maine total	580	43.3	39.8-46.9	28.4	26.0 - 30.7
Sex					
Female	416	60.9		33.2	29.9 - 36.4
Male	164	25.0		20.4	17.3 - 23.5
Age					
Under 5 yrs	*	*	*	NApp	NApp
05-14 yrs	*	*	*	NApp	NApp
15-24 yrs	*	*	*	NApp	NApp
25-34 yrs	*	*	*	NApp	NApp
35-44 yrs	*	*	*	NApp	NApp
45-54 yrs	*	*	*	NApp	NApp
55-64 yrs	*	*	*	NApp	NApp
65-74 yrs	42	25.7	18.0 - 33.5	NApp	NApp
75-84 yrs	166	209.6	177.7 - 241.4	NApp	NApp
85 yrs and over	366	1,095.2	983.0 - 1207.4	NApp	NApp

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.
Alzheimer's Disease Deaths: Deaths with Alzheimer's disease as the underlying cause of death.
Alzheimer's Disease: ICD-10 code G30.
Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
95% CI: 95% confidence interval of the rate.
Subgroup counts might not sum to Maine total due to missing data or rounding.
NApp = Not applicable; age-adjusted rates are not calculated for age-specific groups.
* = To protect privacy, counts less than 10 are suppressed by CDC Wonder.

Table 41. Alzheimer's Disease Deaths by Race and Hispanic Ethnicity, Maine, 2009 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	502	38.0	27.0	26.0 - 27.5
Race				
American Indian or Alaska Native	*	*	*	*
Asian or Pacific Islander	1	^	^	7.8 - 27.9
Black or African American	*	*	*	*
White	499	39.0	26.9	26.1 - 27.6
Hispanic Ethnicity				
Hispanic or Latino	*	*	*	*
Non-Hispanic	500	38.1	26.7	26 - 27.5

Data Source: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov>
 Alzheimer's Disease Deaths: Deaths with Alzheimer's disease as the underlying cause of death.

Alzheimer's Disease: ICD-10 code G30.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

* = To protect privacy, counts less than 10 are suppressed by CDC Wonder.

^: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

Table 42. Alzheimer's Disease Deaths by County, Maine, 2014 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	547	41.0	27.9	26.8 - 28.9
County				
Androscoggin	75	69.6	54.5	49.0 - 60.0
Aroostook	30	43.7	24.3	20.4 - 28.2
Cumberland	111	38.2	27.5	25.2 - 29.8
Franklin	11	35.3	24.3	18.2 - 31.8
Hancock	22	40.7	24.5	19.9 - 29.0
Kennebec	44	36.6	25.9	22.5 - 29.4
Knox	21	52.3	27.9	22.5 - 33.3
Lincoln	13	38.0	19.3	14.9 - 24.6
Oxford	24	41.5	27.8	22.8 - 32.8
Penobscot	42	27.7	20.3	17.5 - 23.0
Piscataquis	9	52.2	29.6	21.5 - 39.7
Sagadahoc	7	20.4	13.7	9.6 - 18.6
Somerset	15	29.9	22.3	17.6 - 27.9
Waldo	12	30.4	22.5	17.2 - 29.0
Washington	10	31.7	19.7	14.6 - 25.9
York	101	49.7	33.5	30.6 - 36.5

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Alzheimer's Disease Deaths: Deaths with Alzheimer's disease as the underlying cause of death.

Alzheimer's Disease: ICD-10 code G30.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 43. Diabetes Deaths by Year, Maine and U.S., 1999-2018

Year	Maine			U.S.	
	Number of Deaths	Age-adjusted Rate Rate	95% CI	Age-adjusted Rate Rate	95% CI
1999	348	24.5	21.9 - 27.0	25.0	24.8 - 25.2
2000	357	24.8	22.2 - 27.4	25.0	24.9 - 25.2
2001	398	27.1	24.4 - 29.7	25.4	25.2 - 25.5
2002	404	27.1	24.4 - 29.7	25.6	25.4 - 25.8
2003	398	26.2	23.6 - 28.8	25.5	25.3 - 25.7
2004	382	24.7	22.2 - 27.2	24.7	24.6 - 24.9
2005	385	24.7	22.2 - 27.1	24.9	24.8 - 25.1
2006	343	21.4	19.1 - 23.7	23.6	23.4 - 23.7
2007	355	21.8	19.5 - 24.1	22.8	22.6 - 22.9
2008	344	20.8	18.6 - 23.0	22.0	21.9 - 22.2
2009	350	20.6	18.5 - 22.8	21.0	20.9 - 21.2
2010	376	21.8	19.6 - 24.1	20.8	20.7 - 21.0
2011	353	20.0	17.9 - 22.1	21.6	21.5 - 21.8
2012	380	21.3	19.1 - 23.5	21.2	21.0 - 21.3
2013	373	20.4	18.3 - 22.5	21.2	21.0 - 21.3
2014	414	22.4	20.2 - 24.6	20.9	20.8 - 21.1
2015	407	21.9	19.7 - 24.1	21.3	21.1 - 21.4
2016	463	23.9	21.7 - 26.2	21.0	20.9 - 21.2
2017	395	19.7	17.7 - 21.7	21.5	21.3 - 21.6
2018	479	23.4	21.3 - 25.66	21.4	21.2 - 21.5

Maine 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. Data Source for years 1999-2017 (ME + US) and US 2018 Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death. 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Feb 13, 2020

Diabetes Deaths: Deaths with diabetes as the underlying cause of death.

Diabetes: ICD-10 codes E10-E14.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 44. Diabetes Deaths by Sex and Year, Maine, 1999-2018						
Year	Maine Males			Maine Females		
	Number of Deaths	Age-adjusted Rate	95% CI	Number of Deaths	Age-adjusted Rate	95% CI
1999	153	26.7	22.4 - 31.0	195	22.6	19.4 - 25.8
2000	155	26.6	22.4 - 30.9	202	23.6	20.3 - 26.9
2001	186	31.3	26.7 - 35.9	212	24.3	21.0 - 27.6
2002	191	30.7	26.3 - 35.2	213	24.2	20.9 - 27.5
2003	198	31.6	27.1 - 36.0	200	22.1	19.0 - 25.2
2004	170	26.3	22.3 - 30.4	212	23.7	20.5 - 26.9
2005	192	29.6	25.4 - 33.9	193	21.3	18.3 - 24.4
2006	161	23.6	19.9 - 27.3	182	19.4	16.6 - 22.3
2007	189	27.2	23.3 - 31.2	166	17.4	14.7 - 20.1
2008	195	27.9	23.9 - 31.9	149	15.5	13.0 - 18.0
2009	188	25.3	21.6 - 29.0	162	16.7	14.1 - 19.4
2010	212	28.2	24.3 - 32.0	164	16.9	14.2 - 19.5
2011	194	24.9	21.3 - 28.5	159	15.9	13.4 - 18.4
2012	223	28.2	24.4 - 32.0	157	15.9	13.3 - 18.4
2013	213	26.0	22.4 - 29.5	160	15.4	12.9 - 17.8
2014	243	29.6	25.7 - 33.4	171	15.9	13.5 - 18.4
2015	226	27.3	23.6 - 30.9	181	17.6	14.9 - 20.3
2016	266	30.9	27.1 - 34.8	197	17.7	15.2 - 20.3
2017	231	25.6	22.2 - 29.0	166	14.0	11.8 - 16.2
2018	271	29.4	25.8 - 33.0	208	18.1	15.6 - 20.7

Data Source for years 1999-2017: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov> . 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Diabetes Deaths: Deaths with diabetes as the underlying cause of death.

Diabetes: ICD-10 codes E10-E14.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 45. Diabetes Deaths by Sex and Age, Maine, 2018

	Number of Deaths	Crude Rate		Age-Adjusted Rate	
		Rate	95% CI	Rate	95% CI
Maine total	479	35.78	32.5-38.9	23.4	21.3 - 25.6
Sex					
Female	208	30.5		18.1	15.6 - 20.7
Male	2711	41.24		29.3	25.7 - 32.9
Age					
Under 5 yrs	*	*	*	NApp	NApp
05-14 yrs	*	*	*	NApp	NApp
15-24 yrs	*	*	*	NApp	NApp
25-34 yrs	*	*	*	NApp	NApp
35-44 yrs	*	*	*	NApp	NApp
45-54 yrs	29	16.1	10.2-22.0	NApp	NApp
55-64 yrs	85	40.1	31.6-48.6	NApp	NApp
65-74 yrs	134	82.0	68.1-95.9	NApp	NApp
75-84 yrs	131	164.1	135.9-192.3	NApp	NApp
85 yrs and over	95	284.3	227.1-341.4	NApp	NApp

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Diabetes Deaths: Deaths with diabetes as the underlying cause of death.

Diabetes: ICD-10 codes E10-E14.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

NApp = Not applicable; age-adjusted rates are not calculated for age-specific groups.

DNR: Data not reliable; death rates are suppressed as unreliable when the numerator is 20 or less.

* = To protect privacy, counts less than 10 are suppressed.

Table 46. Diabetes Deaths by Race and Hispanic Ethnicity, Maine, 2009 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	399	30.0	21.6	20.9 - 22.3
Race				
American Indian or Alaska Native	4	34.9	50.7	34.0 - 72.9
Asian or Pacific Islander	2	^	^	7.7 - 24.6
Black or African American	2	8.8	17.5	10.0 - 28.3
White	392	30.6	21.5	20.8 - 22.2
Hispanic Ethnicity				
Hispanic or Latino	2	10.2	20.4	12.3 - 31.8
Non-Hispanic	395	30.1	21.5	20.8 - 22.2

Data Source: Detailed Mortality Files accessed through CDC Wonder.

<https://wonder.cdc.gov>

Diabetes Deaths: Deaths with diabetes as the underlying cause of death.

Diabetes: ICD-10 codes E10-E14.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

^: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

* = To protect privacy, counts less than 10 are suppressed by CDC Wonder.

Table 47. Diabetes Deaths by County, Maine, 2014 - 2018				
	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	430	32.3	22.2	21.3 - 23.2
County				
Androscoggin	34	31.3	24.2	20.4 - 27.9
Aroostook	29	42.5	25.2	21.0 - 29.3
Cumberland	64	22.0	16.2	14.4 - 18.0
Franklin	14	47.3	34.0	26.2 - 43.4
Hancock	18	33.4	21.1	16.8 - 26.0
Kennebec	46	37.8	27.3	23.7 - 31.0
Knox	13	32.2	18.7	14.3 - 24.0
Lincoln	12	36.3	20.9	15.6 - 27.4
Oxford	30	52.3	35.1	29.4 - 40.9
Penobscot	51	33.7	24.2	21.2 - 27.2
Piscataquis	8	46.2	25.5	18.0 - 35.2
Sagadahoc	7	18.7	12.6	8.5 - 18.0
Somerset	19	37.0	24.9	20.0 - 30.6
Waldo	12	29.9	20.2	15.2 - 26.3
Washington	11	34.8	19.8	14.9 - 25.8
York	63	31.0	21.1	18.7 - 23.5
<p>Data Source: Maine CDC, Data, Research, and Vital Statistics. December 2019.</p> <p>Diabetes Deaths: Deaths with diabetes as the underlying cause of death.</p> <p>Diabetes: ICD-10 codes E10-E14.</p> <p>Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.</p> <p>95% CI: 95% confidence interval of the rate.</p> <p>Subgroup counts might not sum to Maine total due to missing data or rounding.</p> <p>DNR: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.</p>				

Table 48. Influenza and Pneumonia Deaths by Year, Maine and U.S, 1999-2018

Year	Maine			U.S.	
	Number of Deaths	Age-adjusted Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
1999	287	19.9	17.6 - 22.2	23.5	23.3 - 23.7
2000	336	22.9	20.5 - 25.4	23.7	23.6 - 23.9
2001	269	18.0	15.8 - 20.1	22.2	22.0 - 22.4
2002	317	20.9	18.6 - 23.2	23.2	23.0 - 23.3
2003	327	21.4	19.1 - 23.7	22.6	22.4 - 22.7
2004	308	20.0	17.8 - 22.3	20.4	20.2 - 20.5
2005	352	22.3	20.0 - 24.7	21.0	20.9 - 21.2
2006	269	16.7	14.7 - 18.8	18.4	18.3 - 18.6
2007	236	14.2	12.3 - 16.0	16.8	16.7 - 17.0
2008	263	15.5	13.7 - 17.4	17.6	17.5 - 17.7
2009	251	14.6	12.8 - 16.4	16.5	16.4 - 16.7
2010	234	13.4	11.7 - 15.1	15.1	15.0 - 15.2
2011	302	16.8	14.9 - 18.7	15.7	15.6 - 15.9
2012	205	11.6	10.0 - 13.2	14.4	14.3 - 14.6
2013	258	14.0	12.3 - 15.8	15.9	15.7 - 16.0
2014	258	13.7	12.0 - 15.4	15.1	15.0 - 15.3
2015	333	17.6	15.7 - 19.6	15.2	15.1 - 15.4
2016	231	12.0	10.4 - 13.6	13.5	13.4 - 13.6
2017	301	15.2	13.4 - 16.9	14.3	14.2 - 14.4
2018	311	15.2	13.5 - 16.9	14.9	14.8 - 15.0

Maine 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. Data Source for years 1999-2017 (ME + US) and US 2018 Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death. 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Feb 13, 2020

Influenza and Pneumonia Deaths: Deaths with influenza or pneumonia as the underlying cause of death.

Influenza and Pneumonia: ICD-10 codes J10-J18.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 49. Influenza and Pneumonia Deaths by Sex and Year, Maine, 1999-2018

Year	Maine Males			Maine Females		
	Number of Deaths	Age-adjusted Rate	95% CI	Number of Deaths	Age-adjusted Rate	95% CI
1999	118	23.6	19.3 - 28.0	169	17.2	14.6 - 19.9
2000	126	25.7	21.1 - 30.3	210	21.7	18.7 - 24.7
2001	110	21.0	17.0 - 25.0	159	16.3	13.8 - 18.9
2002	141	27.3	22.7 - 31.8	176	17.5	14.9 - 20.1
2003	129	23.7	19.5 - 27.8	198	19.6	16.8 - 22.3
2004	136	23.7	19.7 - 27.7	172	17.6	14.9 - 20.3
2005	148	25.8	21.6 - 30.1	204	20.0	17.2 - 22.7
2006	109	18.5	15.0 - 22.0	160	15.8	13.3 - 18.3
2007	93	15.4	12.4 - 18.9	143	13.5	11.3 - 15.8
2008	112	17.9	14.6 - 21.3	151	14.1	11.8 - 16.3
2009	107	16.1	13.0 - 19.1	144	13.3	11.1 - 15.5
2010	109	16.3	13.2 - 19.4	125	11.4	9.4 - 13.5
2011	140	20.1	16.7 - 23.5	162	14.3	12.1 - 16.6
2012	97	13.8	11.2 - 16.9	108	10.3	8.3 - 12.4
2013	111	15.2	12.4 - 18.1	147	12.9	10.8 - 15.1
2014	112	14.2	11.5 - 16.9	146	12.9	10.7 - 15.0
2015	138	18.3	15.2 - 21.4	195	17.0	14.5 - 19.4
2016	117	14.5	11.8 - 17.2	114	10.1	8.2 - 12.0
2017	135	16.7	13.8 - 19.5	166	14.0	11.8 - 16.2
2018	168	20.1	16.9 - 23.2	143	11.7	9.8 - 13.7

Data Source for years 1999-2017: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov> . 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.
 Influenza and Pneumonia Deaths: Deaths with influenza or pneumonia as the underlying cause of death.
 Influenza and Pneumonia: ICD-10 codes J10-J18.
 Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.

Table 50. Influenza and Pneumonia Deaths by Sex and Age, Maine, 2018

	Number of Deaths	Crude Rate		Age-Adjusted Rate	
		Rate	95% CI	Rate	95% CI
Maine total	311	23.2	20.7-25.8	15.2	13.5 - 16.9
Sex					
Female	143	20.9		11.7	9.8 - 13.7
Male	168	25.6		20.1	16.9 - 23.2
Age					
Under 5 yrs	*	*	*	NApp	NApp
05-14 yrs	*	*	*	NApp	NApp
15-24 yrs	*	*	*	NApp	NApp
25-34 yrs	*	*	*	NApp	NApp
35-44 yrs	*	*	*	NApp	NApp
45-54 yrs	*	*	*	NApp	NApp
55-64 yrs	31	14.6	9.5-19.8	NApp	NApp
65-74 yrs	55	33.7	24.8-42.6	NApp	NApp
75-84 yrs	70	88.4	67.7-109.1	NApp	NApp
85 yrs and over	142	424.9	355.0-494.8	NApp	NApp

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Influenza and Pneumonia Deaths: Deaths with influenza or pneumonia as the underlying cause of death.

Influenza and Pneumonia: ICD-10 codes J10-J18.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

NApp = Not applicable; age-adjusted rates are not calculated for age-specific groups.

* = To protect privacy, counts less than 10 are suppressed

Table 51. Influenza and Pneumonia Deaths by Race and Hispanic Ethnicity, Maine, 2009 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	269	20.2	14.4	13.9 - 15.0
Race				
American Indian or Alaska Native	*	*	*	*
Asian or Pacific Islander	*	*	*	*
Black or African American	*	*	*	*
White	267	20.8	14.5	13.9 - 15.1
Hispanic Ethnicity				
Hispanic or Latino	*	*	*	*
Non-Hispanic	267	20.4	14.4	13.9 - 15.0

Data Source: Detailed Mortality Files accessed through CDC Wonder.
<https://wonder.cdc.gov>

Influenza and Pneumonia Deaths: Deaths with influenza or pneumonia as the underlying cause. Influenza and Pneumonia: ICD-10 codes J09-J18.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

* = To protect privacy, counts less than 10 are suppressed by CDC Wonder.

^ = Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

**Table 52. Influenza and
Pneumonia Deaths by County, Maine, 2014 - 2018**

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	286	21.5	14.7	13.9 - 15.5
County				
Androscoggin	25	22.7	17.3	14.2 - 20.4
Aroostook	20	29.3	17.0	13.8 - 20.8
Cumberland	54	18.7	13.4	11.8 - 15.1
Franklin	8	25.3	19.1	13.3 - 26.6
Hancock	13	23.1	14.0	10.6 - 18.1
Kennebec	24	19.5	14.1	11.5 - 16.6
Knox	7	18.6	10.6	7.2 - 14.9
Lincoln	8	22.2	11.6	8.1 - 16.1
Oxford	12	21.3	14.1	10.8 - 18.2
Penobscot	37	24.6	17.9	15.3 - 20.4
Piscataquis	4	23.7	14.1	8.2 - 22.6
Sagadahoc	8	22.7	14.6	10.5 - 19.9
Somerset	9	18.5	13.8	10.0 - 18.4
Waldo	8	20.3	14.9	10.6 - 20.3
Washington	12	36.7	21.7	16.3 - 28.2
York	38	18.6	12.9	11.1 - 14.8

Data Source: Maine CDC, Data, Research, and Vital Statistics. March 2020.

Influenza and Pneumonia Deaths: Deaths with influenza or pneumonia as the underlying cause of death.

Influenza and Pneumonia: ICD-10 codes J10-J18.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 53. Suicide Deaths by Year, Maine and U.S, 1999-2018

Year	Maine			U.S.	
	Number of Deaths	Age-adjusted Rate Rate	Age-adjusted Rate 95% CI	Age-adjusted Rate Rate	Age-adjusted Rate 95% CI
1999	175	13.4	11.4 - 15.4	10.5	10.3 - 10.6
2000	154	11.8	10.0 - 13.7	10.4	10.3 - 10.5
2001	161	12.1	10.2 - 14.0	10.7	10.6 - 10.8
2002	166	12.5	10.6 - 14.4	10.9	10.8 - 11.1
2003	137	10.0	8.3 - 11.7	10.8	10.7 - 10.9
2004	171	12.4	10.5 - 14.3	11.0	10.9 - 11.1
2005	175	12.4	10.6 - 14.3	10.9	10.8 - 11.0
2006	158	11.0	9.3 - 12.8	11.0	10.9 - 11.1
2007	191	13.6	11.7 - 15.6	11.3	11.2 - 11.4
2008	181	12.7	10.8 - 14.6	11.6	11.5 - 11.7
2009	197	14.0	12.0 - 16.0	11.8	11.6 - 11.9
2010	186	13.2	11.2 - 15.1	12.1	12.0 - 12.2
2011	235	16.6	14.4 - 18.8	12.3	12.2 - 12.5
2012	209	14.5	12.5 - 16.6	12.6	12.5 - 12.7
2013	245	17.4	15.1 - 19.7	12.6	12.5 - 12.7
2014	220	15.7	13.5 - 17.8	13.0	12.9 - 13.1
2015	235	16.0	13.9 - 18.2	13.3	13.2 - 13.4
2016	226	15.9	13.7 - 18.1	13.5	13.4 - 13.6
2017	274	18.9	16.5 - 21.2	14.0	13.9 - 14.1
2018	269	18.4	16.1 - 20.7	14.2	14.1 - 14.3

Maine 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. Data Source for years 1999-2017 (ME + US) and US 2018 Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death. 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Feb 13, 2020

Suicide Deaths: Deaths with suicide as the underlying cause of death.

Suicide: ICD-10 codes X60-X84, Y87.0 .

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 54. Suicide Deaths by Sex and Year, Maine, 1999-2018

Year	Maine Males			Maine Females		
	Number of Deaths	Age-adjusted Rate Rate	Age-adjusted Rate 95% CI	Number of Deaths	Age-adjusted Rate Rate	Age-adjusted Rate 95% CI
1999	137	22.5	18.7 - 26.3	38	5.7	4.0 - 7.8
2000	134	21.5	17.9 - 25.2	20	3.2	1.9 - 4.9
2001	128	20.3	16.8 - 23.9	33	4.8	3.3 - 6.8
2002	133	21.2	17.6 - 24.8	33	4.8	3.3 - 6.8
2003	108	16.7	13.5 - 19.9	29	4.1	2.7 - 5.9
2004	137	20.8	17.3 - 24.3	34	4.9	3.4 - 6.8
2005	145	21.9	18.3 - 25.5	30	4.2	2.8 - 6.0
2006	126	18.2	15.0 - 21.4	32	4.6	3.1 - 6.5
2007	156	23.3	19.5 - 27.0	35	4.9	3.3 - 6.8
2008	146	21.6	18.0 - 25.2	35	4.7	3.2 - 6.6
2009	155	22.7	19.0 - 26.3	42	5.8	4.1 - 8.0
2010	151	21.9	18.4 - 25.5	35	5.1	3.5 - 7.2
2011	185	27.4	23.3 - 31.4	50	7.0	5.1 - 9.3
2012	169	24.2	20.4 - 28.0	40	5.5	3.9 - 7.6
2013	193	28.4	24.3 - 32.6	52	7.2	5.3 - 9.6
2014	174	25.2	21.3 - 29.1	46	6.7	4.9 - 9.1
2015	186	26.9	22.8 - 30.9	49	5.9	4.3 - 8.0
2016	172	24.3	20.5 - 28.1	54	7.9	5.8 - 10.4
2017	216	31.3	27.0 - 35.7	58	7.4	5.5 - 9.7
2018	217	30.4	26.1 - 34.6	52	7.1	5.2 - 9.5

Data Source for years 1999-2017: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov> . 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Suicide Deaths: Deaths with suicide as the underlying cause of death.

Suicide: ICD-10 codes X60-X84, Y87.0.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 55. Suicide Deaths by Sex and Age, Maine, 2018

	Number of Deaths	Crude Rate		Age-Adjusted Rate	
		Rate	95% CI	Rate	95% CI
Maine total	269	20.1	17.7 - 22.5	18.4	16.1 - 20.7
Sex					
Female	52	7.6		7.1	5.2 - 9.5
Male	217	33.1		30.4	26.1 - 34.6
Age					
Under 5 yrs	*	*	*	NApp	NApp
05-14 yrs	*	*	*	NApp	NApp
15-24 yrs	29	19.0	12.1 - 26.0	NApp	NApp
25-34 yrs	36	22.5	15.1 - 29.8	NApp	NApp
35-44 yrs	34	22.3	14.8 - 29.8	NApp	NApp
45-54 yrs	48	26.6	19.1 - 34.2	NApp	NApp
55-64 yrs	56	26.4	19.5 - 33.3	NApp	NApp
65-74 yrs	35	21.4	14.3 - 28.5	NApp	NApp
75-84 yrs	21	26.5	15.2 - 37.8	NApp	NApp
85 yrs and over	*	*	*	NApp	NApp

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Suicide Deaths: Deaths with suicide as the underlying cause of death.

Suicide: ICD-10 codes X60-X84, Y87.0.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

NApp = Not applicable; age-adjusted rates are not calculated for age-specific groups.

DNR: Data not reliable; death rates are suppressed as unreliable when the numerator is 20 or less.

* = To protect privacy, counts less than 10 are suppressed

Table 56. Suicide Deaths by Race and Hispanic Ethnicity, Maine, 2009 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	230	17.3	16.1	15.4 - 16.8
Race				
American Indian or Alaska Native	2	19.3	22.8	13.3 - 36.5
Asian or Pacific Islander	1	^	^	3.2 - 11.3
Black or African American	2	10.2	10.0	6.2 - 15.3
White	224	17.5	16.2	15.5 - 16.9
Hispanic Ethnicity				
Hispanic or Latino	1	^	^	3.7 - 12.0
Non-Hispanic	227	17.3	16.1	15.4 - 16.8

Data Source: Detailed Mortality Files accessed through CDC Wonder.
<https://wonder.cdc.gov>

Suicide Deaths: Deaths with suicide as the underlying cause of death.

Suicide: ICD-10 codes X60-X84, Y87.0 .

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

^: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

* = To protect privacy, counts less than 10 are suppressed by CDC Wonder.

Table 57. Suicide Deaths by County, Maine, 2014 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	244	18.3	16.9	15.9 - 17.9
County				
Androscoggin	20	18.8	17.8	14.4 - 21.7
Aroostook	14	19.9	18.7	14.2 - 24.2
Cumberland	38	13.2	12.3	10.5 - 14.1
Franklin	6	20.6	19.0	12.5 - 27.6
Hancock	10	18.7	15.8	11.4 - 21.2
Kennebec	25	20.5	19.3	15.8 - 22.9
Knox	8	19.6	17.9	12.3 - 25.1
Lincoln	7	20.5	20.2	13.6 - 28.8
Oxford	10	17.8	15.2	11.0 - 20.4
Penobscot	27	18.0	16.9	13.9 - 19.8
Piscataquis	4	23.7	23.2	13.5 - 37.1
Sagadahoc	9	24.4	23.2	16.4 - 31.8
Somerset	12	23.2	20.9	15.6 - 27.4
Waldo	9	22.3	19.6	13.9 - 26.8
Washington	9	27.2	26.0	18.2 - 36.0
York	36	17.9	16.8	14.2 - 19.4

Data Source: Maine CDC. Data, Research, Vital Statistics. December 2019

Suicide Deaths: Deaths with suicide as the underlying cause of death.

Suicide: ICD-10 codes X60-X84, Y87.0 .

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 58. Nephritis, Nephrotic Syndrome & Nephrosis Deaths by Year, Maine and U.S, 1999-2018

Year	Maine			U.S.	
	Number of Deaths	Age-adjusted Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
1999	191	13.4	11.5 - 15.2	13.0	12.9 - 13.2
2000	175	12.0	10.2 - 13.8	13.5	13.4 - 13.6
2001	208	14.0	12.1 - 15.9	14.1	13.9 - 14.2
2002	232	15.4	13.4 - 17.4	14.4	14.3 - 14.5
2003	265	17.4	15.3 - 19.4	14.7	14.5 - 14.8
2004	264	17.1	15.0 - 19.1	14.5	14.3 - 14.6
2005	250	15.9	13.9 - 17.8	14.7	14.5 - 14.8
2006	252	15.7	13.8 - 17.7	14.8	14.7 - 15.0
2007	269	16.5	14.5 - 18.5	14.9	14.8 - 15.0
2008	263	15.6	13.7 - 17.5	15.1	15.0 - 15.3
2009	272	16.3	14.3 - 18.2	15.1	14.9 - 15.2
2010	272	15.8	13.9 - 17.7	15.3	15.2 - 15.4
2011	257	14.8	12.9 - 16.6	13.4	13.3 - 13.5
2012	257	14.3	12.5 - 16.1	13.1	13.0 - 13.2
2013	252	13.6	11.9 - 15.2	13.2	13.1 - 13.3
2014	223	11.7	10.1 - 13.3	13.2	13.1 - 13.3
2015	211	11.4	9.8 - 13.0	13.4	13.3 - 13.5
2016	239	12.4	10.8 - 14.0	13.1	13.0 - 13.3
2017	260	13.0	11.4 - 14.5	13.0	12.9 - 13.1
2018	258	12.5	11.0 - 14.1	12.9	12.8 - 13.0

Maine 2018 data source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. Data Source for years 1999-2017 (ME + US) and US 2018 Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death. 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Feb 13, 2020

Nephritis, Nephrotic Syndrome and Nephrosis Deaths: Deaths with nephritis, nephrotic syndrome or nephrosis as the underlying cause of death.

Nephritis, Nephrotic Syndrome and Nephrosis: ICD-10 codes N00-N07, N17-N19, N25-N27.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 59. Nephritis, Nephrotic Syndrome, & Nephrosis Deaths by Sex and Year, Maine, 1999-2018

Year	Maine Males			Maine Females		
	Number of Deaths	Age-adjusted Rate	95% CI	Number of Deaths	Age-adjusted Rate	95% CI
1999	95	18.2	14.7 - 22.4	96	10.9	8.8 - 13.4
2000	82	15.9	12.6 - 19.8	93	10.0	8.1 - 12.3
2001	97	18.4	14.9 - 22.5	111	11.7	9.5 - 14.0
2002	101	17.9	14.3 - 21.4	131	13.9	11.5 - 16.4
2003	123	21.7	17.8 - 25.6	142	15.2	12.6 - 17.7
2004	104	17.8	14.3 - 21.2	160	17.2	14.5 - 19.9
2005	111	18.8	15.2 - 22.3	139	14.4	12.0 - 16.8
2006	121	19.8	16.2 - 23.3	131	13.4	11.1 - 15.8
2007	135	21.2	17.5 - 24.8	134	13.5	11.2 - 15.8
2008	122	19.0	15.6 - 22.4	141	13.7	11.4 - 16.0
2009	128	19.3	15.9 - 22.7	144	14.1	11.8 - 16.5
2010	146	21.9	18.3 - 25.5	126	12.0	9.9 - 14.2
2011	132	19.0	15.7 - 22.3	125	11.7	9.7 - 13.8
2012	129	17.9	14.8 - 21.0	128	11.8	9.7 - 13.9
2013	134	17.9	14.8 - 21.0	118	10.8	8.8 - 12.7
2014	117	15.4	12.6 - 18.2	106	9.2	7.4 - 11.0
2015	119	15.8	12.9 - 18.7	92	8.3	6.6 - 10.2
2016	113	14.1	11.5 - 16.8	126	11.1	9.1 - 13.1
2017	135	16.1	13.3 - 18.8	125	10.8	8.9 - 12.7
2018	143	16.8	14.0 - 19.6	115	9.6	7.8 - 11.3

Data Source for years 1999-2017: Detailed Mortality Files accessed through CDC Wonder. <https://wonder.cdc.gov> . 2018 data source: Maine Center for Disease

Control and Prevention. Data, Research and Vital Statistics.

Nephritis, Nephrotic Syndrome and Nephrosis Deaths: Deaths with nephritis, nephrotic syndrome or nephrosis as the underlying cause of death.

Nephritis, Nephrotic Syndrome and Nephrosis: ICD-10 codes N00-N07, N17-N19, N25-N27.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Table 60. Nephritis, Nephrotic Syndrome, & Nephrosis Deaths by Sex and Age, Maine, 2018

	Number of Deaths	Crude Rate		Age-Adjusted Rate	
		Rate	95% CI	Rate	95% CI
Maine total	258	19.3	16.9 - 21.6	12.5	11.0 - 14.1
Sex					
Female	115	16.8		9.6	7.8 - 11.3
Male	143	21.8		16.8	14.0 - 19.6
Age					
Under 5 yrs	*	*	*	NApp	NApp
05-14 yrs	*	*	*	NApp	NApp
15-24 yrs	*	*	*	NApp	NApp
25-34 yrs	*	*	*	NApp	NApp
35-44 yrs	*	*	*	NApp	NApp
45-54 yrs	*	*	*	NApp	NApp
55-64 yrs	27	12.7	7.9 - 17.5	NApp	NApp
65-74 yrs	48	29.4	21.1 - 37.7	NApp	NApp
75-84 yrs	67	84.6	64.3 - 104.8	NApp	NApp
85 yrs and over	108	323.2	262.2 - 384.1	NApp	NApp

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Nephritis, Nephrotic Syndrome and Nephrosis Deaths: Deaths with nephritis, nephrotic syndrome or nephrosis as the underlying cause of death.

Nephritis, Nephrotic Syndrome and Nephrosis: ICD-10 codes N00-N07, N17-N19, N25-N27.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

NApp = Not applicable; age-adjusted rates are not calculated for age-specific groups.

DNR: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

* = To protect privacy, counts less than 10 are suppressed by CDC Wonder.

Table 61. Nephritis, Nephrotic Syndrome & Nephrosis Deaths by Race and Hispanic Ethnicity, Maine, 2009 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	250	18.8	13.5	13.0 - 14.0
Race				
American Indian or Alaska Native	1	^	^	6.7 - 34.5
Asian or Pacific Islander	*	*	*	*
Black or African American	1	^	^	4.5 - 20.3
White	247	19.3	13.5	13.0 - 14.1
Hispanic Ethnicity				
Hispanic or Latino	*	*	*	*
Non-Hispanic	249	19.0	13.5	12.9 - 14.0

Data Source: Detailed Mortality Files accessed through CDC Wonder.

<https://wonder.cdc.gov>

Nephritis, Nephrotic Syndrome and Nephrosis Deaths: Deaths with nephritis, nephrotic syndrome or nephrosis as the underlying cause of death.

Nephritis, Nephrotic Syndrome and Nephrosis: ICD-10 codes N00-N07, N17-N19, N25-N27.

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

^: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.

* = To protect privacy, counts less than 10 are suppressed by CDC Wonder.

Table 62. Nephritis, Nephrotic Syndrome, & Nephrosis Deaths by County, Maine, 2014 - 2018

	Average Annual Number	Crude Rate	Age-Adjusted Rate	
			Rate	95% CI
Maine total	238	17.9	12.2	11.5 - 12.9
County				
Androscoggin	12	11.5	8.7	6.6 - 11.2
Aroostook	18	26.4	15.3	12.3 - 18.9
Cumberland	48	16.5	12.1	10.5 - 13.6
Franklin	7	22.0	14.9	10.3 - 21.0
Hancock	12	21.6	12.8	9.7 - 16.5
Kennebec	19	15.8	11.1	9.0 - 13.6
Knox	7	18.1	9.8	6.8 - 13.6
Lincoln	5	14.0	7.6	4.7 - 11.6
Oxford	12	20.6	12.9	9.8 - 16.7
Penobscot	34	22.1	16.2	13.7 - 18.6
Piscataquis	3	15.4	8.4	4.3 - 14.7
Sagadahoc	7	21.0	13.7	9.7 - 18.9
Somerset	6	11.8	8.1	5.4 - 11.6
Waldo	5	13.7	10.6	6.9 - 15.7
Washington	7	22.2	14.1	9.6 - 19.9
York	36	17.8	12.3	10.5 - 14.2

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.

Nephritis, Nephrotic Syndrome and Nephrosis Deaths: Deaths with nephritis, nephrotic syndrome or nephrosis as the underlying cause of death.

Nephritis, Nephrotic Syndrome and Nephrosis: ICD-10 codes N00-N07, N17-N19, N25-N27. Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

Subgroup counts might not sum to Maine total due to missing data or rounding.

Technical Notes

Cause of Death and ICD 10 codes

Causes (as presented in this report)	ICD-10 Codes	Technical Name	Notes
Cancer	C00-C97	Malignant neoplasms	
Heart Disease	I00-I09, I11, I13, I20-I51	Diseases of the heart	
Chronic Lower Respiratory Disease	J40-J47	Chronic Lower Respiratory Disease	Includes chronic obstructive pulmonary disease and asthma
Unintentional Injury	V01-X59, Y85-86	Unintentional Injury	Includes (but not limited to) unintentional drug overdose, motor vehicle crash deaths, and falls
Cerebrovascular Disease	I60-I69	Cerebrovascular disease	Also known as stroke.
Alzheimer's Disease	G30, G309	Alzheimer's Disease	
Diabetes	E10-E14	Diabetes mellitus	Includes both Type 1 and Type 2 diabetes deaths.
Influenza and Pneumonia	J09-J18	Influenza and Pneumonia	
Suicide	X60-X84, Y87.0	Intentional self-harm (suicide)	ICD codes related to terrorism are preceded with '*'
Nephritis, Nephrotic Syndrome and Nephrosis	N00-N07, N17-N19, N25-N27	Nephritis, Nephrotic Syndrome and Nephrosis	Specific disease of the kidneys
Chronic Liver Disease and Cirrhosis	K70, K73-K74	Chronic Liver Disease and Cirrhosis	
Parkinson's Disease	G20-G21	Parkinson's Disease	
Septicemia	A40-A41	Septicemia	A serious infection of the bloodstream (sometimes called blood poisoning or bacteremia) causing sepsis, a life-threatening immune response.
In Situ Neoplasms, Benign Neoplasms and Neoplasms of Uncertain or Unknown Behavior	D00-D48	In Situ Neoplasms, Benign Neoplasms and Neoplasms of Uncertain or Unknown Behavior	Includes (but not limited to) brain and central nervous system tumors and myelodysplastic syndromes.
Pneumonitis due to Solids and Liquids	J69	Pneumonitis due to Solids and Liquids	Pneumonitis (inflammation of lungs) due to inhalation of solids or liquids.

Data Sources:

- Data on leading causes of death by age group and years of potential life lost (YPLL) were obtained from the National Center for Health Statistics (NCHS), National Vital Statistics System. Centers for Disease Control and Prevention: WISQARS Leading Causes of Death. Available at: www.cdc.gov/injury/wisqars
- Mortality data for the US and Maine for years prior to 2018 are from the Centers for Disease Control, National Center for Health Statistics, Underlying Cause of Death. 1999-2018 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999-2018, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> in February 2020.
- Maine mortality data for 2018 are from the Maine CDC, Office of Data, Research and Vital Statistics.
- Counts and rates for Maine may differ slightly from WONDER due to revisions to death records after final year-end counts were submitted to NCHS/WONDER.

Trend Analysis:

- The trend analysis referenced in the Key Summary section as well as in the At-a-Glance table is based on joinpoint regression analysis of WONDER data 1999-2018. Joinpoint is a software developed by CDC that allows one to test changes in trend over a time period and the significance of a given change. If a trend has changed, joinpoint allows multiple segments to be fitted to a graph and to test the significance of a given segment (increasing, decreasing or not significantly different from 0). Joinpoint also calculates the Average Annual Percent Change (AAPC) which assesses how much trends change over time. If a joinpoint identified a significant increase or decrease in trend, the progress was noted with red or green circle, respectively. If the analysis noted that the rate had plateaued (not changed from zero), then the progress was noted as a yellow circle.

References


Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Web-based Injury Statistics Query and Reporting System (WISQARS) [online]. (2017). Available at: www.cdc.gov/injury/wisqars

Joinpoint Regression Program, Version 4.7.0.0 - February 2019 Statistical Methodology and Applications Branch, Surveillance Research Program, National Cancer Institute.

Mokdad AH, Marks JS, Stroup DF, Geberding JL. Actual causes of death in the United States, 2000. *JAMA*. 2004; 291(10):1238–45 (and 2005 erratum).

Office of Data, [Research and Vital Statistics](#), Maine Center for Disease Control and Prevention, Department of Health and Human Services.

United States Department of Health and Human Services (US DHHS), Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death 1999-2018 on CDC WONDER Online Database, released February 13, 2020. Data are from the Multiple Cause of Death Files, 1999- 2018 as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html>



The Department of Health and Human Services (DHHS) does not discriminate on the basis of disability, race, color, creed, gender, sexual orientation, age, or national origin, in admission to, access to, or operations of its programs, services, or activities, or its hiring or employment practices. This notice is provided as required by Title II of the Americans with Disabilities Act of 1990 and in accordance with the Civil Rights Act of 1964 as amended, Section 504 of the Rehabilitation Act of 1973, as amended, the Age Discrimination Act of 1975, Title IX of the Education Amendments of 1972 and the Maine Human Rights Act and Executive Order Regarding State of Maine Contracts for Services. Questions, concerns, complaints or requests for additional information regarding the ADA may be forwarded to DHHS' ADA Compliance/EEO Coordinators, 11 State House Station – 221 State Street, Augusta, Maine 04333, 207-287-4289 (V), 207-287-3488 (V), TTY users call Maine relay 711. Individuals who need auxiliary aids for effective communication in program and services of DHHS are invited to make their needs and preferences known to the ADA Compliance/EEO Coordinators. This notice is available in alternate formats, upon request.